

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
5 July 2001 (05.07.2001)

PCT

(10) International Publication Number  
**WO 01/48183 A2**

- (51) International Patent Classification<sup>7</sup>: **C12N 15/00**
- (21) International Application Number: **PCT/EP00/13149**
- (22) International Filing Date:  
22 December 2000 (22.12.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
9930691.2 24 December 1999 (24.12.1999) GB
- (71) Applicant (for all designated States except US): **DEV-GEN NV** [BE/BE]; Technologiepark 9, B-9052 Zwijnaarde (BE).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **PLAETINCK, Geert** [BE/BE]; Pontstraat 16, B-9820 Merelbeke (BE). **MORTIER, Katherine** [BE/BE]; Paddenhoek 20, B-9830 St.-Martens Latem (BE). **LISSENS, Ann** [BE/BE]; Tiensesteenweg 137, B-3010 Kessel-Lo (BE). **BOGAERT, Thierry** [BE/BE]; Wolvendreef 26g, B-8500 Kortrijk (BE).
- (74) Agent: **BAYLISS, Geoffrey, Cyril**; Boulton Wade Tennant, Verulam Gardens, 70 Gray's Inn Road, London WC1X 8BT (GB).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**  
— Without international search report and to be republished upon receipt of that report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



**WO 01/48183 A2**

(54) Title: IMPROVEMENTS RELATING TO DOUBLE-STRANDED RNA INHIBITION

(57) Abstract: There are described ways of improving the efficiency of double stranded RNA inhibition as a method of inhibiting gene expression in nematode worms such as *C. elegans*. In particular, the invention relates to the finding that changes in the genetic background of *C. elegans* result in increased sensitivity to double-stranded RNA inhibition.

IMPROVEMENTS RELATING TO DOUBLE-STRANDED RNAINHIBITION

5 The present invention is concerned with ways of  
improving the efficiency of double stranded RNA  
inhibition as a method of inhibiting gene expression  
in nematode worms such as *C. elegans*. In particular,  
the invention relates to the finding that the  
susceptibility of nematode worms such as *C. elegans* to  
10 double stranded RNA inhibition is affected by changes  
in the genetic background of the worms.

It has recently been described in Nature Vol 391,  
pp.806-811, February 98, that introducing double  
stranded RNA into a cell results in potent and  
15 specific interference with expression of endogenous  
genes in the cell, which interference is substantially  
more effective than providing either RNA strand  
individually as proposed in antisense technology. This  
specific reduction of the activity  
20 of the gene was also found to occur in the nematode  
worm *Caenorhabditis elegans* (*C. elegans*) when the RNA  
was introduced into the genome or body cavity of the  
worm.

The present inventors have utilized the double  
25 stranded RNA inhibition technique and applied it  
further to devise novel and inventive methods of (i)  
assigning functions to genes or DNA fragments which  
have been sequenced in various projects, such as, for  
example, the human genome project and which have yet  
30 to be accorded a particular function, and (ii)  
identifying DNA responsible for conferring a  
particular phenotype. Such methods are described in  
the applicant's co-pending application number WO  
00/01846. Processes for introducing RNA into a living  
35 cell, either *in vivo* or *ex vivo*, in order to inhibit  
expression of a target gene in that cell are

additionally described in WO 99/32619.

Several different experimental approaches can be used to introduce double-stranded RNA into nematode worms in order to achieve RNA interference *in vivo*.

5 One of the most straightforward approaches is simple injection of double-stranded RNA into a body cavity. A more elegant solution is to feed the nematodes on food organisms, generally bacteria, which express a double stranded RNA of the appropriate sequence,  
10 corresponding to a region of the target gene.

The present inventors have now determined that the phenomenon of RNA interference in nematodes following ingestion of food organisms capable of expressing double-stranded RNA is dependent both on  
15 the nature of the food organism and on the genetic background of the nematodes themselves. These findings may be exploited to provided improved methods of double-stranded RNA inhibition.

Therefore, according to a first aspect of the  
20 present invention there is provided a method of inhibiting expression of a target gene in a nematode worm comprising feeding to said nematode worm a food organism which is capable of producing a double-stranded RNA structure having a nucleotide sequence  
25 substantially identical to a portion of said target gene following ingestion of the food organism by the nematode, wherein the nematode has a non wild-type genetic background selected to provide increased sensitivity to RNA interference as compared to wild  
30 type.

*Caenorhabditis elegans* is the preferred nematode worm for use in the method of the invention although the method could be carried out with other nematodes and in particular with other microscopic nematodes,  
35 preferably microscopic nematodes belonging to the genus *Caenorhabditis*. As used herein the term "microscopic" nematode encompasses nematodes of

approximately the same size as *C. elegans*, being of the order 1mm long in the adult stage. Microscopic nematodes of this approximate size can easily be grown in the wells of a multi-well plate of the type  
5 generally used in the art to perform mid- to high-throughput screening.

It is an essential feature of this aspect of the invention that the nematode has a non wild-type genetic background which confers greater sensitivity  
10 to RNA interference phenomena (abbreviated herein to RNAi) as compared to the equivalent wild type nematodes. As illustrated in the accompanying examples, introduction of double-stranded RNA (abbreviated herein to dsRNA) into a non wild-type  
15 strain according to the invention results in greater inhibition of expression of the target gene. Depending on the nature of the target gene, this greater level of inhibition may be detectable at the phenotypic level as a more pronounced phenotype.

20 The nematode having non wild-type genetic background may, advantageously, be a mutant strain. Mutations which have the effect of increasing susceptibility of the nematode to RNAi may, for example, affect the stability of dsRNA or the kinetics  
25 of dsRNA turnover within cells of the worm or the rate of uptake of dsRNA synthesised by a food organism. Suitable mutant strains include mutant strains exhibiting knock-out or loss-of-function mutations in one or more genes encoding proteins involved in RNA  
30 synthesis, RNA degradation or the regulation of these processes.

In one preferred embodiment, the nematode is a mutant strain, more preferably a mutant *C. elegans*, which exhibits reduced activity of one or more  
35 nucleases compared to wild-type. Suitable strains include mutant strains exhibiting knock-out or loss-of-function mutations in one or more genes encoding

nucleases, such as RNases. A particularly preferred example is the *nuc-1* strain. This mutant *C. elegans* strain is known per se in the art.

5 In a second preferred embodiment, the nematode is a mutant strain, more preferably a mutant *C. elegans*, which exhibits increased gut uptake compared to wild-type. Particularly preferred examples of such strains are the so-called *C. elegans* gun mutants described herein. In a still further embodiment, the nematode  
10 may be a transgenic worm comprising one or more transgenes which increase gut uptake relative to wild-type.

The term "increased gut uptake" as used herein is taken to mean increased uptake of foreign particles  
15 from the gut lumen and may encompass both increased gut permeability and increased gut molecular transport compared to wild-type *C. elegans*.

*C. elegans* feeds by taking in liquid containing its food (e.g. bacteria). It then spits out the  
20 liquid, crushes the food particles and internalises them into the gut lumen. This process is performed by the muscles of the pharynx. The process of taking up liquid and subsequently spitting it out is called pharyngeal pumping. Once the food particles have been  
25 internalised via pharyngeal pumping their contents must cross the gut itself in order to reach target sites in the worm. There are multiple factors which effect the uptake of compounds from the gut lumen to the surrounding tissues. These include the action of  
30 multi-drug resistance proteins, multi-drug resistance related proteins and the P450 cytochromes as well as other enzymes and mechanisms available for transport of molecules through the gut wall.

*C. elegans* mutants which exhibit increased uptake  
35 of foreign molecules through the gut may be obtained from the *C. elegans* mutant collection at the C.

*C. elegans* Genetic Center, University of Minnesota, St Paul, Minnesota, or may be generated by standard methods. Such methods are described by Anderson in Methods in Cell Biology, Vol 48, "C. elegans: Modern biological analysis of an organism" Pages 31 to 58. Several selection rounds of the PCR technique can be performed to select a mutant worm with a deletion in a desired gene. Alternatively, a population of worms could be subjected to random mutagenesis and worms exhibiting the desired characteristic of increased gut uptake selected using a phenotypic screen, such as the dye uptake method described herein.

As an alternative to mutation, transgenic worms may be generated with the appropriate characteristics. Methods of preparing transgenic worms are well known in the art and are particularly described by Craig Mello and Andrew Fire, Methods in Cell Biology, Vol 48, Ed. H.F. Epstein and D.C. Shakes, Academic Press, pages 452-480.

Worms exhibiting the desired characteristics of increased gut uptake can be identified using a test devised by the inventors based on uptake of a marker precursor molecule which is cleaved by the action of enzymes present in the gut lumen to generate a marker molecule which produces a detectable signal, such as fluorescence. A suitable marker precursor molecule is the fluorescent dye precursor BCECF-AM available from Molecular Probes (Europe BV), Netherlands. This dye only becomes fluorescent when cleaved by esterases and maintained at a pH above 6. The pH of the gut lumen is usually 5 or below. Thus, any BCECF-AM taken up through the pharynx into the gut lumen is not fluorescent until cleaved and the cleaved portion has entered the cells surrounding the lumen which are at a higher pH. Thus, this dye is able to quickly identify mutant or otherwise modified worms which have increased gut transport or permeability. There is a

gradual increase in fluorescence in the tissues surrounding the gut while the gut lumen remains dark. The fluorescence can be detected at an excitation wavelength of 485 nm and an emission wavelength of 530 nm.

Specific examples of gun mutant strains isolated using this procedure which may be used in the method of the invention are strains bg77, bg84, bg85 and bg86, although it is to be understood that the invention is in no way limited to the use of these specific strains. The *C. elegans* gun mutant strain bg85 was deposited on 23 December 1999 at the BCCM/LMG culture collection, Laboratorium Voor Microbiologie, Universiteit Gent, K. L. Ledeganckstraat 35, B-9000, Gent, Belgium under accession number LMBP 5334CB. The phrase "the bg85 mutation" as used herein refers to the specific mutation(s) present in the bg85 strain which is/are responsible for conferring the gun phenotype.

It is also within the scope of the invention to use a non wild-type nematode strain, preferable a *C. elegans* strain, having multiple mutations which affect sensitivity to RNAi. A preferred type of multiple mutant is one having at least one mutation which results in reduced nuclease activity compared to wild type and at least one mutation which results in increased gut uptake compared to wild type. An example of such a mutant is a *C. elegans* strain having the *nuc-1* mutation and at least one further gun mutation. As exemplified herein, double mutants having the *nuc-1* mutation and a gun mutation exhibit enhanced sensitivity to RNAi as compared to either *nuc-1* or gun single mutants.

For the avoidance of doubt, where particular characteristics or properties of nematode worms are described herein by relative terms such as "enhanced"

or "increased" or "decreased" this should be taken to mean enhanced, increased or decreased relative to wild-type nematodes. In the case of *C. elegans*, wild-type is defined as the N2 Bristol strain which is well known to workers in the *C. elegans* field and has been extremely well characterised (see Methods in Cell Biology, Volume 48, *Caenorhabditis elegans: Modern biological analysis of an organism*, ed. by Henry F. Epstein and Diane C. Shakes, 1995 Academic Press; The nematode *Caenorhabditis elegans*, ed. by William Wood and the community of *C. elegans* researchers., 1988, Cold Spring Harbor Laboratory Press; *C. elegans* II, ed. by Donald L. Riddle, Thomas Blumenthal, Barbara J. Meyer and James R. Priess, 1997, Cold Spring Harbor Laboratory Press). The N2 strain can be obtained from the *C. elegans* Genetic Center, University of Minnesota, St Paul, Minnesota, USA.

The food organism for use in the above aspect of the invention is preferably a bacterium such as, for example, a strain of *E.coli*. It will, however, be appreciated that any other type of food organism on which nematodes feed and which is capable of producing dsRNA could be used. The food organism may be genetically modified to express a double-stranded RNA of the appropriate sequence, as will be understood with reference to the examples included herein. One convenient way in which this may be achieved in a bacterial food organism is by transforming the bacterium with a vector comprising a promoter or promoters positioned to drive transcription of a DNA sequence to RNA capable of forming a double-stranded structure. Examples of such vectors will be further described below.

The actual step of feeding the food organism to the nematode may be carried out according to procedures known in the art, see WO 00/01846.



Typically the feeding of the food organisms to the nematodes is performed on standard agar plates commonly used for culturing *C. elegans* in the laboratory. However, the step of feeding the food  
5 organism to the nematodes may also be carried out in liquid culture, for example in the wells of 96-well microtitre assay plates.

The inventors have further observed that variations in the food organism can result in enhanced  
10 *in vivo* RNAi when the food organism is ingested by a nematode worm.

Accordingly, in a further aspect the invention provides a method of inhibiting expression of a target gene in a nematode worm comprising feeding to said  
15 nematode worm a food organism capable of producing a double-stranded RNA structure having a nucleotide sequence substantially identical to a portion of said target gene following ingestion of the food organism by the nematode, wherein the food organism carries a  
20 modification selected to provide increased expression or persistence of the double-stranded RNA compared to a food organism which does not carry the modification.

The modification present in the food organism can be any modification which results in increased  
25 expression of the dsRNA or in increased persistence of the dsRNA. Suitable modifications might include mutations within the bacterial chromosome which affect RNA stability and/or degradation or mutations which have a direct effect on the rate of transcription. In  
30 a preferred embodiment, the food organism is an RNase III minus *E. coli* strain, or any other RNase negative strain.

According to a still further aspect of the invention there is provided a method of inhibiting  
35 expression of a target gene in a nematode worm comprising introduction of a DNA capable of producing a double-stranded RNA structure having a nucleotide

13 sequence substantially identical to a portion of said target gene in said nematode, wherein the nematode is one which exhibits increased gut uptake compared to wild type.

5 In addition to exhibiting increased sensitivity to RNAi following feeding with food organisms capable of expressing a dsRNA, nematodes which exhibit increase gut uptake as described herein also show increased uptake of DNA molecules capable of producing  
10 double-stranded RNA structures following ingestion into a nematode.

In a preferred embodiment, the DNA is in the form of a vector comprising a promoter or promoters orientated to relative to a sequence of DNA such that  
15 they are capable of driving transcription of the said DNA to make RNA capable of forming a double-stranded structure upon binding of an appropriate RNA polymerase to the promoter or promoters.

Several different arrangements of promoters may  
20 be used in such a vector. In a first arrangement a DNA fragment corresponding to a region of the target gene is flanked by two opposable polymerase-specific promoters which are preferably identical. Transcription from the opposable promoters produces  
25 two complementary RNA strands which can anneal to form an RNA duplex. The plasmid pGN1 described herein is an example of a vector comprising two opposable T7 promoters flanking a multiple cloning site for insertion of a DNA fragment of the appropriate  
30 sequence, corresponding to a region of a target gene. pGN8 is an example of a vector derived from pGN1 containing a fragment of the *C. elegans unc-22* gene. In an alternative arrangement, DNA fragments corresponding to a region of the target gene may be  
35 placed both in the sense and the antisense orientation downstream of a single promoter. In this case, the sense/antisense fragments are co-transcribed to

generate a single RNA strand which is self-complementary and can therefore form an RNA duplex.

In both of the above arrangements, the polymerase-specific T3, T7 and SP6 promoters, all of which are well known in the art, are useful for driving transcription of the RNA. Expression from these promoters is dependent on expression of the cognate polymerase. Advantageously, the nematode itself may be adapted to express the appropriate polymerase. Expression of the polymerase may be general and constitutive, but could also be regulated under a tissue-specific promoter, an inducible promoter, a temporally regulated promoter or a promoter having a combination of such characteristics. Transgenic *C. elegans* strains harboring a transgene encoding the desired polymerase under the control of an appropriately-regulated promoter can be constructed according to methods known *per se* in the art and described, for example, by Craig Mello and Andrew Fire in *Methods in Cell Biology*, Vol 48, Ed. H. F. Epstein and D. C. Shakes, Academic Press, pp 452-480.

The advantage of adapting the nematode to express the required polymerase is that it is possible to control inhibition of expression of the target gene in a tissue-specific and/or temporally specific manner by placing expression of the polymerase under the control of an appropriately regulated promoter.

Introduction of DNA into nematodes in accordance with the method of the invention can be achieved using a variety of techniques, for example by direct injection into a body cavity or by soaking the worms in a solution containing the DNA. If the DNA is in the form of a vector as described herein, e.g. a plasmid harboring a cloned DNA fragment between two flanking T7 promoters, then dsRNA corresponding to this DNA fragment will be formed in the nematode resulting in down regulation of the corresponding gene. The

introduced DNA can form an extrachromosomal array,  
which array might result in a more catalytic knock-out  
or reduction of function phenotype. The DNA might also  
become integrated into the genome of the nematode,  
5 resulting in the same catalytic knock out or reduction  
of function phenotype, but which is stably  
transmittable.

In each aspect of the invention, the double-  
stranded RNA structure may be formed by two separate  
10 complementary RNA strands or a single self-  
complementary strand, as described above. Inhibition  
of target gene expression is sequence-specific in that  
only nucleotide sequences corresponding to the duplex  
region of the dsRNA structure are targeted for  
15 inhibition.

It is preferred to use dsRNA comprising a  
nucleotide sequence identical to a portion of the  
target gene, although RNA sequences with minor  
variations such as insertions, deletions and single  
20 base substitutions may also be used and are effective  
for inhibition. It will be readily apparent that 100%  
sequence identity between the dsRNA and a portion of  
the target gene is not absolutely required for  
inhibition and the phrase "substantially identical" as  
25 used herein is to be interpreted accordingly.  
Generally sequences which are substantially identical  
will share at least 90%, preferably at least 95% and  
more preferably at least 98% nucleic acid sequence  
identity. Sequence identity may be conveniently  
30 calculated based on an optimal alignment, for example  
using the BLAST program accessible at  
WWW.ncbi.nlm.nih.gov.

The invention will be further understood with  
reference to the following non-limiting Examples,  
35 together with the accompanying Figures in which:

Figure 1 is a plasmid map of the vector pGN1

containing opposable T7 promoters flanking a multiple cloning site and an ampicillin resistance marker.

5 Figure 2 is a plasmid map of the vector pGN8 (a genomic fragment of the *C. elegans unc-22* gene cloned in pGN1).

10 Figure 3 is a plasmid map of the vector pGN29 containing two T7 promoters and two T7 terminators flanking *Bst*XI sites. This vector permits cloning of DNA fragments linked to *Bst*XI adaptors.

15 Figure 4 is a plasmid map of the vector pGN39 containing two T7 promoters and two T7 terminators flanking attR recombination sites (based on the Gateway™ cloning system of Life Technologies, Inc).

20 Figure 5 is a plasmid map of the vector pGX22 (a fragment of the *C. elegans* gene C04H5.6 cloned in pGN29).

25 Figure 6 is a plasmid map of the vector pGX52 (a fragment of the *C. elegans* gene K11D9.2b cloned in pGN29).

Figure 7 is a plasmid map of the vector pGX104 (a fragment of the *C. elegans* gene Y57G11C.15 cloned in pGN29).

30 Figure 8 is a plasmid map of the vector pGZ8 (a fragment of the *C. elegans* gene T25G3.2 cloned in pGN39).

35 Figure 9 shows the results of an RNAi experiment in which wild-type (N2) or *nuc-1* strain *C. elegans* in liquid culture were fed with *E. coli* containing the

plasmid pGX22.

Figure 10 shows the results of an RNAi experiment in which wild-type (N2) or *nuc-1* strain *C. elegans* in liquid culture were fed with *E. coli* containing the plasmid pGX52.

Figure 11 shows the results of an RNAi experiment in which wild-type (N2) or *nuc-1* strain *C. elegans* in liquid culture were fed with *E. coli* containing the plasmid pGXGZ8.

Figure 12 shows the results of an RNAi experiment in which wild-type (N2) or *nuc-1* strain *C. elegans* in liquid culture were fed with *E. coli* containing the plasmid pGX104

**Example 1**

Influence of genetic background on the efficiency of RNAi in *C. elegans*.

**5 Introduction**

Various different *C. elegans* strains were fed with different bacteria, to test the possibility of RNAi by feeding *C. elegans* with bacteria that produce dsRNA. The possibility of DNA delivery and dsRNA delivery has  
 10 previously been envisaged by using different bacterial strains. In this experiment the importance of the *C. elegans* strain as receptor of the dsRNA is also shown.

For this experiment the following *E. coli* strains were  
 15 used:

1. MC1061: F-*araD139*  $\Delta$ (*ara-leu*)7696 *galE15 galk16*  $\Delta$ (*lac*)X74 *rps1* (*Str<sup>r</sup>*) *hsdR2* (*r<sub>k</sub><sup>-</sup> m<sub>k</sub><sup>+</sup>*) *mcrA mcrB1*  
 - regular host for various plasmids,  
 - Wertman et al., (1986) Gene 49:253-262,  
 - Raleigh et al., (1989) in Current Protocols in  
 Molecular Biology eds. Ausubel et al, Publishing  
 associates and Wiley Interscience; New York. Unit  
 1.4.  
 25
2. B21(DE3): F- *ompT(lon)* *hsdS<sub>B</sub>* (*r<sub>B</sub><sup>-</sup>, m<sub>B</sub><sup>-</sup>*; an *E. coli* B  
 strain) with DE3, a  $\lambda$  prophage carrying the T7  
 RNA polymerase gene.  
 - regular host for IPTG inducible T7 polymerase  
 expression,  
 - Studier et al. (1990) Meth. Enzymol. 185:60-89  
 30
3. HT115 (DE3): F- *mcrA mcrb* IN(*rrnD-rrnE*) 1  $\lambda$ -  
*rnc14::tr10* (DE3 lysogen: *lacUV5*  
 promoter-T7polymerase)  
 35 - host for IPTG inducible T7 polymerase

expression,  
- RNaseIII-,  
- Fire A, Carnegie Institution, Baltimore, MD,  
Pers. Comm.

5

For this experiment the following *C. elegans* strains were used:

1. *C. elegans* N2: regular WT laboratory strain
2. *C. elegans nuc-1(e1393)*: *C. elegans* strain with a reduced endonuclease activity (>95%); condensed chromatin persists after programmed cell death; ingested (bacterial) DNA in the intestinal lumen is not degraded. Several alleles are described:  
e1392 (strong allele: has been used for the experiments described below); n887 (resembles e1392) and n334 (weaker allele)  
- Stanfield et al. (1998) East Coast Worm meeting abstract 171,  
- Anonymous, Worm Breeder's Gazette 1(1):17b  
Hevelone et al. (1988) Biochem. Genet. 26:447-461  
- Ellis et al., Worm breeder's Gazette 7(2):44  
- Babu, Worm Breeder's gazette 1(2):10  
- Driscoll, (1996) Brain Pathol. 6:411-425  
- Ellis et al., (1991) Genetics 129:79-94

10

15

20

25

For this experiment the following plasmids were used:

pGN1: A vector encoding for ampicillin resistance, harbouring a multiple cloning site between two convergent T7 promoters.

pGN8: pGN1 containing a genomic fragment of *unc-22*.  
Decreased *unc-22* expression via RNAi results in a "twitching" phenotype in *C. elegans*.

35



**Experimental conditions**

12-well micro-titer plates were filled with approximately 2 ml of NGM agar per well (1 litre of NGM agar: 15g Agar, 1g peptone, 3g NaCl, 1ml cholesterol solution (5 mg/ml in EtOH), with sterile addition after autoclaving of 9.5 ml 0.1M CaCl<sub>2</sub>, 9.5 ml 0.1 ml MgSO<sub>4</sub>, 25 ml 1M KH<sub>2</sub>PO<sub>4</sub>/K<sub>2</sub>HPO<sub>4</sub> buffer pH 6 and 5 ml nystatin solution (dissolved 10 mg/ml in 1:1 EtOH:CH<sub>3</sub>COONH<sub>4</sub> 7.5 M).

The dried plates were spotted with approximately 50 µl of an overnight culture of bacteria. When IPTG induction was required, 50 µl of a 10 mM stock solution of IPTG was dropped on top of the bacteria lawn, and incubated at 37°C for approximately 4 hours. Individual nematodes at the L4 growth stage were then placed in single wells. In each well 4 nematodes, and the plates were further incubated at 20°C for 6 days to allow offspring to be formed. The F1 offspring of the seeded nematodes were tested for the twitching phenotype.

## Results

Table 1: Percentage of the offspring that show the twitching phenotype

5	<b>MC1061</b>	<b>N2</b>	<b><i>nuc-1</i></b>
	pGN1	0%	0%
	pGN1 + IPTG	0%	0%
	pGN8	0%	0%
	pGN8 + IPTG	0%	0%
10	<b>BL21 (DE3)</b>		
	pGN1	0%	0%
	pGN1 + IPTG	0%	0%
	pGN8	20% (+)	>90% (++)
	pGN8 + IPTG	20% (+)	>90% (++±)
15	<b>HT115 (DE3)</b>		
	pGN1	0%	0%
	pGN1 + IPTG	0%	0%
	pGN8	50% (±)	>90% (++)
	pGN8 + IPTG	80% (++)	>90% (+++)

5: %: percentage twitchers

+: weak twitching

++: twitching

+++: strong twitching

## Conclusions

The experiment with *E. coli* MC1061 shows that no twitching could be observed in this experiment.

Neither the N2 nematodes nor the *nuc-1* nematodes showed any twitchers. This is to be expected as *E. coli* MC1061 does not produce any T7 RNA polymerase, and hence the *unc-22* fragment cloned in pGN8 is not

expressed as dsRNA.

5 The experiment with *E. coli* strain BL21(DE3) and  
nematode strain N2 shows expected results. BL21(DE3)  
harbouring plasmid pGN1 does not result in any  
twitching as the pGN1 vector is an empty vector. BL21  
(DE3) harbouring PGN8 results in the expression of  
unc-22 dsRNA. When this dsRNA is fed to the N2  
10 nematode (indirectly by feeding with the bacteria that  
produce the dsRNA), this results in a twitching  
phenotype, indicating that the dsRNA is able to pass  
the gut barrier and is able to perform its interfering  
activity.

15 Surprisingly the RNAi effect of the unc-22 dsRNA was  
even more pronounced in *C. elegans* strain *nuc-1* than  
in the wild type N2 strain. Although one may expect  
that the *nuc-1* mutation results in the non-degradation  
or at least in a slower degradation of DNA, as the  
20 NUC-1 protein is known to be involved in DNase  
activity, we clearly observe an enhancement of the  
RNAi induced phenotype in *C. elegans* with a *nuc-1*  
background. The *nuc-1* mutation has not been cloned  
yet, but it has been described that the gene is  
25 involved in nuclease activity, and more particularly  
DNase activity. If the NUC-1 protein is a nuclease, it  
may also have activity on nuclease activity on dsRNA,  
which would explain the enhanced RNAi phenotype. The  
*nuc-1* gene product may be a nuclease, or a regulator  
30 of nuclease activity. As the mode of action of RNAi is  
still not understood, it is also possible that the  
NUC-1 protein is interfering in the mode of action of  
RNAi. This would explain why a *nuc-1* mutant is more  
sensitive to RNAi.

35

The experiment with the *E.coli* strain HT115 (DE3)

confirms the experiments with the BL21(DE3) strain. The RNA interference observed with the unc-22 dsRNA is even higher. In comparison with strain BL21(DE3) this could be expected, as HT115(DE3) is a RNase III minus strain, and hence is expected to produce larger amounts of dsRNA, resulting in more prominent RNAi. This indicates further that the RNAi observed in this experiment is the result of the dsRNA produced by the bacteria fed to the *C. elegans*. Feeding *C. elegans* nuc-1 with HT115(DE3) harbouring pGN8 also results in higher RNA interference phenotype than feeding the same bacteria to *C. elegans* wild-type strain N2. Once again this indicates that improved RNAi can realised using a nuclease negative *C. elegans* and more particularly with a with the *C. elegans* nuc-1 (e1392) strain.

### Summary

RNA interference can be achieved in *C. elegans* by feeding the worms with bacteria that produce dsRNA. The efficiency of this RNA interference is dependent both on the *E. coli* strain and on the genetic background of the *C. elegans* strain. The higher the level of dsRNA production in the *E. coli*, the more RNAi is observed. This can be realised by using efficient RNA expression systems such as T7 RNA polymerase and RNAase negative strains, such as RNaseIII minus stains. In this example the level of dsRNA production varied: HT115(DE3)>BL21(DE3)>MC1061. RNA interference is high in *C. elegans* strains that are nuclease negative, or that are influenced in their nuclease activity. This can be realised by using a mutant strain such as *C. elegans* nuc-1. In this example the sensitivity to RNAi varied: *C. elegans* nuc-1 >> *C. elegans* N2

**Example 2**

Improved RNAi by feeding dsRNA producing bacteria in selected *C. elegans* strains-Comparison of the *nuc-1* strain with several mutants which show improved gut uptake (designated herein 'gun' mutants). Strains bg77, bg78, bg83, bg84, bg85, bg86, bg87, bg88 and bg89 are typical gun mutant *C. elegans* strains isolated using selection for increased gut uptake (gun phenotype) with the marker dye BCECF-AM.

**Experimental conditions:**

- 12-well micro-titer plates were filled with approximately 2ml of NGM agar (containing 1ml/l of ampicillin (100µg/ml) and 5 ml of 100mM stock IPTG) per well
- the dried plates were spotted with 25µl of an overnight culture of bacteria (BL21DE3/HT115DE3) containing the plasmids pGN1 (T7prom-T7prom) or pGN8 (T7prom-unc-22-T7prom)
- individual nematodes at the L4 growth stage were then placed in single wells, one nematode per well
- the plates were incubated at 20°C for 6 days to allow offspring to be formed
- the adult F1 offspring of the seeded nematodes were tested for the twitching phenotype

**Results:**

Table 2:

	20°C/6d	pGN1 HT115DE3	pGN8 BL2DE3	pGN8 HT115DE3	
5	N2	0	1	1	
	<i>nuc-1</i>	0	1-2	3	
	bg77	0	1-2	3	
	bg78	0	1	1-2	
	bg83	0	1	1	
10	bg84	0	1-2	3	
	bg85	0	1	2-3	
	bg86	0	1	2	
	bg87	0	1	1	
	bg88	0	1	1	
15	bg89	0	1	1	

figure legend:

- 0 = no twitching
- 1 = no to weak phenotype
- 2 = clear phenotype
- 3 = strong phenotype

**Conclusions**

- bacterial strain HT115(DE3) shows a better RNAi sensitivity than bacterial strain BL21(DE3)
- the *nuc-1* *C. elegans* strain is a better strain than the Wild-type N2 strain for RNAi sensitivity
- various gun mutants (improved gut uptake mutants) and more particularly the gun mutant strains bg77, bg84, bg85, bg86 show improved sensitivity to RNAi compared to Wild-type.

**A double mutant *C. elegans* strain (nuc-1/gun) shows even greater sensitivity to RNAi compared to wild-type:**

5 Double mutants were constructed to test the prediction  
that gun/nuc mutants would even show more enhanced  
RNAi sensitivity. As an example, the crossing  
strategy with gun strain bg85 is shown, similar  
crosses can be conducted with other gun strains, such  
10 as bg77, bg84 and bg86.

P0 cross: gun(bg85) x WT males

15 F1 cross: nuc-1 x gun(bg85)/+ males

F2 cross: nuc-1 x gun(bg85)/+; nuc-1/0 males (50%)  
nuc-1 x +/+; nuc-1/0 males (50%)

20 F3 single: gun(bg85)/+; nuc-1 hermaphrodites (25%)  
+/+; nuc-1 hermaphrodites (75%)

F4 single: gun(bg85); nuc-1 (1/4 of every 4th  
plate high staining with BCECF)

25 F5 retest: gun(bg85); nuc-1 (100% progeny of F4  
singled high staining with BCECF)

To select for the gun phenotype, the fluorescence  
precursor BCECF-AM is used (obtainable from Molecular  
30 probes). The precursor BCECF-AM is cleaved by  
esterases present in the gut of the worm to generate  
the dye BCECF which is fluorescent at pH values above  
6. This allows selection for worms that have a gun  
phenotype. BCECF-AM is taken up through the pharynx  
35 into the gut lumen and is not fluorescent until it has  
been cleaved, and the BCECF portion has entered the

cells surrounding the lumen. Wild-type worms will show slower or no increase in BCECF fluorescence.

### 5     **Example 3**

Improved RNAi feeding in liquid culture using *nuc-1(e1393)* *C. elegans*.

#### **Introduction**

10     N2 and *nuc-1* *C. elegans* strains were fed with bacteria producing dsRNAs that give lethal phenotypes via RNAi. For this example RNAi was performed in liquid culture instead of on agar plates. We show here for a number of genes that the RNAi effect is more penetrant using  
15     the *nuc-1* strain than the N2 strain, and that RNAi can be performed in liquid.

For this experiment the following *E. coli* strains were used:

20

1.     HT115 (DE3): F- *mcrA mcrb IN(rrnD-rrnE)* 1  $\lambda$ -  
         *rnc14::tr10* (DE3 lysogen: lacUV5 promoter -T7  
         polymerase)  
         - host for IPTG inducible T7 polymerase expression  
25       - RNaseIII-  
         - Fire A, Carnegie Institution, Baltimore, MD,  
         Pers. Comm.

25

For this experiment, following *C. elegans* strains were used:

30

1.     *C. elegans* N2: regular WT laboratory strain
2.     *C. elegans nuc-1(e1393)*: *C. elegans* strain with a  
35       reduced endonuclease activity (>95%); condensed  
         chromatin persists after programmed cell death;

35



- ingested (bacterial) DNA in the intestinal lumen is not degraded. Several alleles are described: e1392 (strong allele: has been used for the experiments described below); n887 (resembles e1392) and n334 (weaker allele)
- 5       - Stanfield et al. (1998) East Coast Worm meeting abstract 171
- Anonymous, Worm Breeder's Gazette 1(1):17b
- Hevelone et al. (1988) Biochem. Genet. 26:447-461
- 10       - Ellis et al., Worm breeder's Gazette 7(2):44
- Babu, Worm Breeder's gazette 1(2):10
- Driscoll, (1996) Brain Pathol. 6:411-425
- Ellis et al., (1991) Genetics 129:79-94

15

For this experiment, the following plasmids that all give lethal phenotypes in *C. elegans* via RNAi were used:

- 20       pGX22: a vector encoding ampicillin resistance, containing a genomic fragment of cosmid C04H5.6 corresponding to a member of the RNA helicase family.

- pGX52: a vector encoding ampicillin resistance, containing a genomic fragment of cosmid K11D9.2b corresponding to sarco/endoplasmic Ca<sup>2+</sup> ATPase also known as SERCA.
- 25

- pGZ18: a vector encoding ampicillin resistance, containing a genomic fragment of cosmid T25G3.2 corresponding to a chitin like synthase gene.
- 30

- pGX104: a vector encoding ampicillin resistance, containing a genomic fragment of cosmid Y57G11C.15 corresponding to sec-61, a transport protein.
- 35

**Experimental conditions**

- 1 ml overnight cultures of HT115 (DE3) bacteria containing the plasmids pGX22, pGX52, pGZ18 or pGX104 respectively were pelleted and resuspended in S-complete medium, containing 1ml/l of ampicillin (100 µg/ml) and 1ml/l of 1000mM IPTG.
  - 10 µl of this bacterial solution was transferred to a 96-well microtiter plate already filled with 100 µl S-complete containing 1ml/l of ampicillin (100 µg/ml) and 1ml/l of 1000mM IPTG.
  - 3 nematodes at the L1 growth stage of N2 and nuc-1 strain were then placed in single wells, 3 L1's per well. Per experimental set up, 16 wells were used (n=16).
  - the plates were incubated at 25°C for 5 days to allow offspring to be formed.
  - the plates were visually checked and the following phenotypes could be scored per individual well:
- no effect:** L1's developed to adults and gave normal offspring.
- no F1 offspring:** L1's developed to adults and gave no offspring.
- acute lethal:** original L1 did not mature and died.

**Results**

- The results of this experiment are illustrated graphically in Figures 9 to 12. Data are expressed as

a percentage of the total (n=16) on the y-axis for both N2 and *nuc-1* strains.

### Conclusions

5 The following genes were tested in this liquid RNAi assay:

- C04H5.6: an RNA helicase. RNAi of this gene interferes with the generation of offspring.
- 10 - SERCA: a sarco/endoplasmic  $\text{Ca}^{2+}$  ATPase. A strong RNAi phenotype causes an acute lethal phenotype. A less penetrant RNAi effect results in loss of offspring.
- T25G3.2: a chitin like synthase gene. RNAi of  
15 this gene causes dead eggs.
- *sec-61*: a transport protein. A strong RNAi phenotype causes an acute lethal phenotype. A less penetrant RNAi effect results in loss of offspring.
- 20 - RNAi can be performed under liquid conditions.

As in the previous examples this set of experiments shows that the *nuc-1 C. elegans* strain is more sensitive to RNAi than the wild-type N2 strain. This  
25 is most clear for less penetrant phenotypes such as SERCA and chitin synthase. For strong RNAi phenotypes like the helicase and *Sec-61* the difference between the N2 wild-type strain and the *nuc-1* stain is less pronounced.

30

**Example 4****Cloning of pGX22, pGX52, pGZ18 and pGX104 for RNAi**

A set of primers for each gene was designed on the basis of sequence data available in the publicly accessible *C. elegans* sequence database (Acedb).

The cosmid names relate to:

1. **C04H5.6**=member of RNA helicase
2. **K11D9.2b**=SERCA
3. **Y57G11C.15**=transport protein sec-61
4. **T25G3.2**=chitin synthase like

The following primer sequences were designed:

1. **C04H5.6F** 5'-TGCTCAGAGAGTTTCTCAACGAACC-3'  
**C04H5.6R** 5'-CAATGTTAGTTGCTAGGACCACCTG-3'
2. **K11D9.2bF** 5'-CAGCCGATCTCCGTCTTGTG-3'  
**K11D9.2bR** 5'-CCGAGGGCAAGACAACGAAG-3'
3. **Y57G11C.15F** 5'-ACCGTGGTACTCTTATGGAGCTCG-3'  
**Y57G11C.15R** 5'-TGCAGTGGATTGGGTCTTCG-3'
4. **T25G3.2F**  
5'-GGGGACAAGTTTGTACAAAAAGCAGGCTATGCCAAGTACATGTCGATTGCG-3'  
**T25G3.2R**  
5'-GGGGACCACTTTGTACAAGAAAGCTGGGTGGAGAAGCATTCCGAGAGTTTG-3'

PCR was performed on genomic DNA of N2 strain *C. elegans* to give PCR products of the following sizes:

- 1326bp for C04H5.6
- 1213bp for K11D9.2b

1024bp for Y57G11C.15

1115bp for T25G3.2

5 The PCR fragments of C04H5.6, K11D9.2b and Y57G11C.15  
were linked to *Bst*XI adaptors (Invitrogen) and then  
cloned into the pGN29 vector cut with *Bst*XI. pGN29  
contains two T7 promoters and two T7 terminators  
flanking a cloning site which is adapted for  
10 facilitated cloning of PCR fragments, comprising a  
stuffer DNA flanked by two *Bst*XI sites (see schematic  
Figure 3). The resulting plasmids were designated  
pGX22 (C04H5.6), pGX52 (K11D9.2b) and pGX104  
(Y57G11C.15).

15 The PCR fragment of T25G3.2 was cloned into pGN39 via  
recombination sites based on the GATEWAY™ cloning  
system (Life Technologies, Inc). pGN39 contains two  
T7 promoters and two T7 terminators flanking a cloning  
site which facilitates "High Throughput" cloning based  
20 on homologous recombination rather than restriction  
enzyme digestion and ligation. As shown schematically  
in Figure 4, the cloning site comprises *att*R1 and  
*att*R2 recombination sites from bacteriophage lambda  
flanking a gene which is lethal to *E. coli*, in this  
25 case the *ccdB* gene. This cloning site is derived from  
the Gateway™ cloning system commercially available  
from Life Technologies, Inc. The Gateway™ cloning  
system has been extensively described by Hartley et  
al. in WO 96/40724 (PCT/US96/10082).

30

**Example 5**

Selecting *C. elegans* mutations for increased gut uptake (gun) using marker dye BCECF-AM and *unc-31* as background.

5

The screen was performed in *unc-31(e928)* mutant background, to ensure high amounts of dye in the gut lumen, since *unc-31* mutations show constitutive pharyngeal pumping. The dye (BCECF-AM: 2',7' bis (2-carboxyethyl)-5-(and-6)-carboxyfluorescein, acetoxymethylester), obtained from Molecular Probes, is cleaved by intracellular esterases. Fluorescence accumulates in the gut cells upon passage through the apical gut membrane.

15

**Mutagenesis**

Day 1: *unc-31* L4 staged worms were mutagenised with EMS (final concentration 50 mM) for 4 hours

Day 2: P0 was divided over several large agar plates

Day 6: F1's were collected and dropped on large plates. The number of eggs the F1's layed were checked every hour and de F1's were removed when 10-20 eggs per F1 were counted

Day 10: F2 adults were collected and screened with BCECF-AM. Mutations with increased staining of the gut cells after 15-30 minutes exposure to the dye were selected and singled on small agar plates.

30

About 50 initial positives gave progeny which was retested with BCECF-AM (2x) and leucine CMB (1x) 9 of the 50 strains were kept (2 strains : 3 times positive, 7 other strains : twice positive)

35

Table 3: Isolation of mutations for increased staining with BCECF-AM

	Total P0	Total F1	Total F2	screened chromosomes	number of strains isolated
5	(counted)	(estimated)	(calculated)	(estimated)	(counted)
	2251	55618	222472	100000	9

Outcrossing, backcrossing and double construction

- 10 1. backcrossing *unc-31; gun* --> *unc-31; gun*
  - *unc-31; gun* x WT males
  - singled 2x5 WT hermaphrodites F1s (*unc-31/+;gun/+*)
  - singled 50 WT hermaphrodites F2s (1/4 homozygous)
  - select strains segregating 1/4 *unc*
- 15 - stain *unc* strains with BCECF-AM
  - from positive strains pick *unc* homozygous
  - retest 100 % *unc* strains with BCECF-AM
  - kept 1 strain (backcrossed)
- 20 2. *unc-31* background was crossed out-->+; *gun*
  - *unc-31; gun* x WT males
  - singled 2x5 WT hermaphrodites F1s (*unc-31/+;gun/+*)
  - singled 50 WT hermaphrodites F2s (1/4 homozygous)
  - select strains which did not segregate *unc* F3s
- 25 anymore
  - stain non *unc* strains with BCECF-AM
  - 7 positive strains were retested with BCECF-AM and finally 1 was selected and kept (outcrossed)
- 30 3. +; *gun* (1x outcrossed) were 2 times backcrossed-->+; *gun* (3x backcrossed)
  - *gun* x WT males
  - WT hermaphrodites x F1 males (*gun/+*)
  - singled 10 WT hermaphrodites F2s (1/2 heterozygous)
- 35 - singled 50 WT hermaphrodites F3s (1/8 homozygous)

- 31 -

- stain strains with BCECF-AM- retested positives with BCECF-AM and finally 1 was selected and kept

4. *gun* (3x backcrossed) were crossed with *nuc-1(X)*  
 5 mutant--> *gun; nuc-1*  
 - *gun* x WT males  
 - *nuc-1* x *gun/+* males  
 - *nuc-1* x *gun/+; nuc-1/0* or *+/+; nuc-1/0* males  
 - singled 10 WT hermaphrodite progeny (*nuc-1*  
 10 homozygous,  $\frac{1}{2}$  heterozygous *gun*)  
 - singled 40 WT hermaphrodite progeny (1/8 homozygous *gun*)  
 - stain strains with BCECF-AM  
 - retested positives with BCECF-AM and finally 1 was  
 15 selected and kept

Table 6: Strains derived from *gun* mutations

20	<i>gun</i>	<i>unc-31; gun</i>		<i>unc-31; gun</i>		+; <i>gun</i>		<i>gun; nuc-1</i>
		original isolate		backcrossed (1x)		outcrossed (1x)		3x b.c.
	allele number	isolation number	strain number	isolation number	strain number	isolation number	strain number	strain number
	bg77	31.4	UG 510	31.4.46.1	UG 556	31.4.34	UG 563	UG 674
25	bg78	37.5	UG 511	37.5.46.4	UG 557	37.5.15	UG 564	UG 675
	bg83	10.2	UG 543	10.2.11	UG 600	10.2.21	UG 586	UG 676
	bg84	7.2	UG 544	7.2.10	UG 601	7.2.15	UG 589	UG 677
	bg85	11.5	UG 545	11.5.29.2	UG 602	2x b.c.	UG 717	
	bg86	42.1	UG 546	42.1.4.5	UG 603	42.1.18	UG 587	UG 678
30	bg87	7.1	UG 547	7.1.8.3	UG 604	7.1.22	UG 585	UG 679
	bg88	5.3	UG 548	5.3.9	UG 605	5.3.18	UG 584	UG 680
	bg89	23.4	UG 549	23.4.13.5	UG 606	23.4.3	UG 588	UG 671

35



SEQUENCE LISTING:

SEQ ID NO: 1 complete sequence of pGN1

5 SEQ ID NO: 2 complete sequence of pGN8

SEQ ID NO: 3 complete sequence of pGN29

SEQ ID NO: 4 complete sequence of pGN39

10 SEQ ID NO: 5 complete sequence of pGX22

SEQ ID NO: 6 complete sequence of pGX52

15 SEQ ID NO: 7 complete sequence of pGX104

SEQ ID NO: 8 complete sequence of pGZ8

SEQ ID NO: 9 primer C04H5.6F

20 SEQ ID NO: 10 primer C04H5.6R

SEQ ID NO: 11 primer K11D9.2bF

25 SEQ ID NO: 12 primer K11D9.2bR

SEQ ID NO: 13 primer Y57G11C.15F

SEQ ID NO: 14 primer Y57G11C.15R

30 SEQ ID NO: 15 primer T25G3.2F

SEQ ID NO: 16 primer T25G3.2R

**Claims:**

1. A method of inhibiting expression of a target gene in a nematode worm comprising feeding to  
5 said nematode worm a food organism which is capable of producing a double-stranded RNA structure having a nucleotide sequence substantially identical to a portion of said target gene following ingestion of the food organism by the nematode, wherein the nematode  
10 has a non wild-type genetic background selected to provide increased sensitivity to RNA interference as compared to wild type.
2. A method as claimed in claim 1 wherein the  
15 nematode is a microscopic nematode.
3. A method as claimed in claim 2 wherein the nematode is from the genus *Caenorhabditis*.
- 20 4. A method as claimed in claim 3 wherein the nematode is *C. elegans*.
5. A method as claimed in any one of claims 1 to 4 wherein the nematode has a mutant genetic  
25 background.
6. A method as claimed in claim 5 wherein the nematode is a mutant strain which exhibits reduced activity of one or more nucleases compared to wild  
30 type.
7. A method as claimed in claim 6 wherein the nematode is *C. elegans* strain *nuc-1*.
- 35 8. A method as claimed in claim 5 wherein the nematode is a mutant strain which exhibits increased

gut uptake compared to wild type.

9. A method as claimed in claim 8 wherein the nematode is mutant *C. elegans* strain bg85.

5

10. A method as claimed in claim 5 wherein the nematode is a mutant strain having at least one mutation which results in reduced nuclease activity compared to wild type and at least one mutation which results in increased gut uptake compared to wild type.

10

11. A method as claimed in claim 10 wherein the nematode is a mutant *C. elegans* strain having the *nuc-1* mutation and the bg85 mutation.

15

12. A method as claimed in any one of the preceding claims wherein the food organism has been engineered to express a double-stranded RNA.

20

13. A method as claimed in any one of the preceding claims wherein the food organism is a bacterium.

25

14. A method as claimed in claim 13 wherein the food organism is *E. coli*.

30

15. A method as claimed in any one of the preceding claims wherein the food organism has been genetically modified to express a double-stranded RNA having a nucleotide sequence substantially identical to a portion of said target gene.

35

16. A method as claimed in claim 15 wherein the food organism contains a DNA vector, the vector comprising a promoter or promoters orientated relative to a DNA sequence such that they are capable of

initiating transcription of said DNA sequence to RNA capable of forming a double-stranded structure upon binding of an appropriate RNA polymerase to said promoter or promoters.

5

17. A method as claimed in claim 25 wherein the vector comprises two promoters flanking the DNA sequence.

10

18. A method as claimed in claim 26 wherein the two promoters are identical.

15

19. A method as claimed in claim 25 wherein the vector comprises a single promoter and further comprises said DNA sequence in a sense and an antisense orientation relative to said promoter.

20

20. A method as claimed in any one of claims 16 to 20 wherein the nematode or the food organism is adapted to express an RNA polymerase capable of initiating transcription from said promoter or promoters.

25

21. A method as claimed in any one of claims 16 to 20 wherein the RNA polymerase is T7, T3 or SP6 polymerase.

30

22. A method as claimed in any one of claims 1 to 21 wherein the step of feeding said food organism to said nematode worm is carried out in liquid culture.

35

23. A method of inhibiting expression of a target gene in a nematode worm comprising feeding to said nematode worm a food organism capable of producing a double-stranded RNA structure having a nucleotide sequence substantially identical to a

portion of said target gene following ingestion of the food organism by the nematode, wherein the food organism carries a modification selected to provide increased expression or persistence of the doubled-stranded RNA compared to a food organism which does not carry the modification.

24. A method as claimed in claim 23 wherein the food organism is a bacterium.

25. A method as claimed in claim 24 wherein the bacterium is an *E. coli* strain.

26. A method as claimed in claim 25 wherein the *E. coli* strain is an RNase III minus strain or any other RNase negative strain.

27. A method as claimed in any one of claims 23 to 26 wherein the step of feeding said food organism to said nematode worm is carried out in liquid culture.

28. A method of inhibiting expression of a target gene in a nematode worm comprising introduction of a DNA capable of producing a double-stranded RNA structure having a nucleotide sequence substantially identical to a portion of said target gene in said nematode, wherein the nematode is one which exhibits increased gut uptake compared to wild type.

29. A method as claimed in claim 28 wherein the nematode is a microscopic nematode.

30. A method as claimed in claim 29 wherein the nematode is from the genus *Caenorhabditis*.

31. A method as claimed in claim 30 wherein the

nematode is *C. elegans*.

32. A method as claimed in any one of claims 28  
to 31 wherein the nematode has a mutant genetic  
5 background.

33. A method as claimed in claim 32 wherein the  
nematode is mutant *C. elegans* strain bg85.

10 34. A method as claimed in any one of claims 28  
to 33 wherein the DNA capable of producing a double-  
stranded RNA structure is a vector comprising a  
promoter or promoters orientated relative to a DNA  
sequence such that they are capable of initiating  
15 transcription of said DNA sequence to RNA capable of  
forming a double-stranded structure upon binding of an  
appropriate RNA polymerase to said promoter or  
promoters.

20 35. A method as claimed in claim 34 wherein the  
vector comprises two promoters flanking the DNA  
sequence.

25 36. A method as claimed in claim 35 wherein the  
two promoters are identical.

30 37. A method as claimed in claim 34 wherein the  
vector comprises a single promoter and further  
comprises said DNA sequence in a sense and an  
antisense orientation relative to said promoter.

35 38. A method as claimed in any one of claims 34  
to 37 wherein the nematode is adapted to express an  
RNA polymerase capable of initiating transcription  
from said promoter or promoters.

39. A method as claimed in any one of claims 34

to 38 wherein the RNA polymerase is T7, T3 or SP6 polymerase.

*FIG. 1.*






FIG. 2.




FIG. 3.




FIG. 4.




FIG. 5.




FIG. 6.




FIG. 7.




FIG. 8.




FIG. 9.




FIG. 10.






FIG. 11.





FIG. 12.



1  
SEQUENCE LISTING

&lt;110&gt; DEVGEN NV

&lt;120&gt; IMPROVEMENTS RELATING TO DOUBLE-STRANDED RNA INHIBITION

&lt;130&gt; SCB/53711/001

&lt;140&gt;

&lt;141&gt;

&lt;160&gt; 14

&lt;170&gt; PatentIn Ver. 2.0

&lt;210&gt; 1

&lt;211&gt; 3216

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Plasmid pGN1

&lt;400&gt; 1

gagtgcacca	tatgcggtgt	gaaataccgc	acagatgcgt	aaggagaaaa	taccgcatca	60
ggcgaaattg	taaacgttaa	tattttggtt	aaattcgcgt	taaatatttg	ttaaatcagc	120
tcatttttta	accaataggc	cgaaatcggc	aaaatccctt	ataaatcaaa	agaatagacc	180
gagataggg	tgagtgttgt	tccagtttgg	aacaagagtc	cactattaaa	gaacgtggac	240
tccaacgtca	aagggcgaaa	aaccgtctat	cagggcgatg	gcccactacg	tgaaccatca	300
cccaaatcaa	gttttttgcg	gtcgaggtgc	cgtaaagctc	taaatcggaa	ccctaaaggg	360
agccccgat	ttagagcttg	acgggggaa	ccggcgaaac	tggcgagaaa	ggaaggggaa	420
aaagcgaaag	gagcgggagc	tagggcgctg	gcaagtgtag	cggtcacgct	gcgcgtaacc	480
accacacccg	ccgcgcttaa	tgcgccgcta	cagggcgctg	ccattcgcca	ttcaggctgc	540
gcaactgttg	ggaagggcga	tcggtgcggg	cctcttcgct	attacgccag	ctggcgaaa	600
ggggatgtgc	tgcaaggcga	ttaagttggg	taacgccagg	gttttcccag	tcacgacgtt	660
gtaaaacgac	ggccagtga	ttgtaatacg	actcactata	gggcgaattc	gagctcggta	720
cccggggatc	ctctagagtc	gaaagcttct	cgccctatag	tgagtcgtat	tacagcttga	780
gtattctata	gtgtcaccta	aatagcttgg	cgtaatcatg	gtcatagctg	tttcctgtgt	840
gaaattgtta	tccgctcaca	attccacaca	acatacgagc	cggaagcata	aagtgtaaag	900
cctgggtgtg	ctaattgagt	agctaactca	cattaattgc	gttgcgctca	ctgccgcgtt	960
tccagtcggg	aaacctgtcg	tgccagctgc	attaatgaat	cgccaacgc	gcggggagag	1020
gcggtttg	tattgggcgc	tcttcgcgtt	cctcgctcac	tgactcgctg	cgctcggtcg	1080
ttcggtcg	gcgagcggta	tcagctcact	caaaggcgg	aatacggtta	tccacagaat	1140
caggggataa	cgcaggaaa	aacatgtgag	caaaaggcca	gcaaaaggcc	aggaaccgta	1200
aaaaggccgc	gttgctggcg	tttttcgata	ggctccgccc	ccctgacgag	catcacaaaa	1260
atcgacgctc	aagtcagagg	tggcgaaaacc	cgacaggact	ataaagatac	caggcgtttc	1320
ccctggaag	ctccctcg	cgctctcctg	ttccgaccct	gccgcttacc	ggataacctgt	1380
ccgcctttct	cccttcggga	agcgtggcgc	tttctcatag	ctcacgctgt	aggtatctca	1440
gttcggtgta	ggtcggtcgc	tccaagctgg	gctgtgtgca	cgaaccccc	gttcagcccg	1500
accgtgcgc	cttatccggt	aactatcgct	ttgagtcctc	cccggtaaga	cacgacttat	1560
cgccactggc	agcagccact	ggtaacagga	ttagcagagc	gaggtatgta	ggcgggtgcta	1620
cagagttctt	gaagtgggtg	cctaactacg	gctacactag	aaggacagta	tttggtatct	1680
gcgctctgct	gaagccagtt	accttcggaa	aaagagttgg	tagctcttga	tccggcaaac	1740
aaaccaccgc	tggtagcgg	ggtttttttg	tttgcaagca	gcagattacg	cgcagaaaaa	1800
aaggatctca	agaagatcct	ttgatctttt	ctacggggtc	tgacgctcag	tggaaacgaaa	1860
actcacgtta	agggattttg	gtcatgagat	tatcaaaaag	gatcttcacc	tagatccttt	1920
taaattaaaa	atgaagtttt	aaatcaatct	aaagtatata	tgagtaaact	tggtctgaca	1980

```

gttaccaatg cttaatcagt gaggcaccta tctcagcgat ctgtctatatt cggttcacca 2040
tagttgcctg actccccgtc gtgtagataa ctacgatacg ggagggctta ccatctggcc 2100
ccagtgtctg aatgataccg cgagaccac gctcaccggc tccagattta tcagcaataa 2160
accagccagc cggaagggcc gagcgagaa gtggctcctgc aactttatcc gcctccatcc 2220
agtctattaa ttgttgccgg gaagctagag taagtagttc gccagttaat agtttgcgca 2280
acgttgttgg cattgctaca ggcacgtgg tgtcacgctc gtcgtttgg atggcttcat 2340
tcagctccgg ttcccaacga tcaaggcgag ttacatgac ccccatgttg tgcaaaaaag 2400
cggttagctc cttcgggtcct ccatcgttg tcagaagtaa gttggccgca gtgttatcac 2460
tcattggttat ggcagcactg cataattctc ttactgtcat gccatccgta agatgctttt 2520
ctgtgactgg tgagtactca accaagtcac tctgagaata ccgcgcccgg cgaccgagtt 2580
gctcttgccc ggcgtcaata cgggataata gtgtatgaca tagcagaact ttaaaagtgc 2640
tcattcattgg aaaacgttct tcggggcgaa aactctcaag gatcttaccg ctgttgagat 2700
ccagttcgat gtaacccact cgtgcaccca actgatcttc agcatctttt actttcacca 2760
gcgtttctgg gtgagcaaaa acaggaaggc aaaatgccgc aaaaaaggga ataagggcga 2820
cacggaaatg ttgaatactc atactcttcc tttttcaata ttattgaagc atttatcagg 2880
gttattgtct catgagcgga tacatatttg aatgtattta gaaaaataaa caaatagggg 2940
ttccgcgcac atttccccga aaagtgccac ctgacgtcta agaaaccatt attatcatga 3000
cattaaccta taaaaatagg cgtatcacga ggccctttcg tctcgcgcgt ttcggtgatg 3060
acggtgaaaa cctctgacac atgcagctcc cgggacgggt cacagcttgt ctgtaagcgg 3120
atgccgggag cagacaagcc cgtcagggcg cgtcagcggg tgttgccggg tgctcggggct 3180
ggcttaacta tgcggcatca gagcagattg tactga 3216

```

&lt;210&gt; 2

&lt;211&gt; 4620

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Plasmid pGN8

&lt;400&gt; 2

```

gatccgaatc tccatgtctg ttaacagcct tgacacggaa tttatattca tgcccttgag 60
tcaaactcgtc aacgtggaag ttggtatcct tgctctctcc gcaagcagtc catctgccag 120
tggcagcatc ttgcttttca atgacatagt gactgatttc agctcctcca tcatcttctg 180
gttccttcca tgcaagatca catccatcct tgacaatatt agtgacatcg agaggtccac 240
gtgggcttga tggatgatca agaacagtaa ccttcacttc agcagtgatc gttccattct 300
cgttctctgc cttgatgata taggttcctg tatccgaacg caaagctctc ttcacatgga 360
atthagtact gccgtcttca ttgttcaact tcatacgatc atcagattcg actggtgttc 420
cttcgaaagt ccaagtaatt gttggagttg gttcaccact gactggaatg ttcaatgaga 480
agtcttgtcc agccttgacc ttgatttctt gaatcgagtt acgatcgatg actggtggaa 540
ctataattta attcaatgat tattagtaat tgatttagac tcttaccatt tctagccttt 600
gcaacagctg atgctgaatc agatggatct cccaatcctg ccttgttctt ggcacggatt 660
ctgaattcgt actttgatcc ttccttgaga tttccaacag tagcattcgt ttgtccagct 720
ggaacatgag caacgtcatt ccagaatggc gagaactcgt ccttcatctc aacaacgat 780
tcctcgattg gagcaccacc gtcgtttgct ggtggcttcc attcaaggtc aacatgatcc 840
ttatcccaat cagtaatttc aggagcattt gtctttcctg gcttgtcaaa tggatctttg 900
gcaagtgtgg ttccgaaggc ctccaatgga tcggactctc cttcagcatt gacggcagcg 960
acacggaact gaaaatcaaa atgttgtagg caattgagtt caagattaaa aaattctcac 1020
tttatattca tgtccaggaa taagaccgtc aacaacagct gtagtcttat ctccagcgac 1080
ctttgcagct ggaacccatc ttccacttgc agtatcgatc ttttcgatca catagttttc 1140
aattggaata cctccatcat catctggtgc acgccaattc aaagtgcacat gatcaccatg 1200
aacatcgga aacatctaag gaccatttgg agaagttggc ttgtctgaaa atttaaaata 1260
taaccaaatt aattgaagaa aactaatgct caccaataac attgatctta acagttgctt 1320
catcttctcc atttgcattg acagctttga tagtgaaagt tccactgtct ccacgttcca 1380
tttgcttca aaccagcttt gattggtatt ctgggttatc aagcttctcg ccctatagtg 1440
agtcgtatta cagcttgagt attctatagt gtcacctaaa tagcttggcg taatcatggg 1500
catagctgtt tcctgtgtga aattgttatc cgctcacaat tccacacaac atacgagccg 1560
gaagcataaa gtgtaaagcc tggggtgcct aatgagtgag ctaactcaca ttaattgcgt 1620

```

tgcgctcact	gcccgcctttc	cagtcgggaa	acctgtcgtg	ccagctgcat	taatgaatcg	1680
gccaacgcgc	ggggagaggc	ggtttgcgta	ttgggcgctc	ttccgcttcc	tcgctcactg	1740
actcgctgcg	ctcggtcgtt	cggctgcggc	gagcggatc	agctcactca	aaggcggtaa	1800
tacggttatc	cacagaatca	ggggataacg	caggaaagaa	catgtgagca	aaaggccagc	1860
aaaaggccag	gaaccgtaaa	aaggccgcgt	tgctggcggt	tttcgatagg	ctccgcccc	1920
ctgacgagca	tcacaaaaat	cgacgctcaa	gtcagagggt	gcgaaacccg	acaggactat	1980
aaagatacca	ggcgctttccc	cctggaagct	ccctcggtcg	ctctcctggt	ccgaccctgc	2040
cgtctaccgg	atacctgtcc	gcctttctcc	cttcgggaag	cgtggcgctt	tctcatagct	2100
cacgctgtag	gtatctcagt	tcgggtgtag	tcgttcgctc	caagctgggc	tgtgtgcacg	2160
aaccccccg	tcagcccgac	cgctgcgcct	tatccggtaa	ctatcgtctt	gagtccaacc	2220
cggtaagaca	cgacttatcg	ccactggcag	cagccactgg	taacaggatt	agcagagcga	2280
ggtagtagg	cgggtgctaca	gagttcttga	agtgggtggc	taactacggc	tacactagaa	2340
ggacagtatt	tggtagctgc	gctctgctga	agccagttac	cttcggaaaa	agagttggta	2400
gctcttgatc	cggcaaacaa	accaccgctg	gtagcgggtg	tttttttgtt	tgcaagcagc	2460
agattacgcg	cagaaaaaaa	ggatctcaag	aagatccttt	gatcttttct	acggggtctg	2520
acgctcagtg	gaacgaaaac	tcacgttaag	ggatttttgt	catgagatta	tcaaaaagga	2580
tcttcacctc	gatcctttta	aattaaaaat	gaagttttaa	atcaatctaa	agtatatatg	2640
agtaaacttg	gtctgacagt	taccaatgct	taatcagtg	ggcacctatc	tcagcgatct	2700
gtctatcttc	ttcatccata	gttgccctg	tccccgtcgt	gtagataaact	acgatacggg	2760
agggccttacc	atctggcccc	agtgtgcaa	tgataccgcg	agaccacgc	tcaccggctc	2820
cagatttatc	agcaataaac	cagccagccg	gaagggccga	gcgcagaagt	ggtcctgcaa	2880
ctttatccgc	ctccatccag	tctattaatt	gttgccggga	agctagagta	agtagttcgc	2940
cagttaatag	tttgcgcaac	gttgttgga	ttgctacagg	catcgtgggt	tcacgctcgt	3000
cgtttgggat	ggcttcattc	agctccggtt	cccaacgatc	aaggcgagtt	acatgatccc	3060
ccatgttgtg	caaaaaagcg	gttagctcct	tcggctcctc	gatcgttgtc	agaagtaagt	3120
tggccgcagt	gttatcactc	atggttatgg	cagcactgca	taattctctt	actgtcatgc	3180
catccgtaag	atgcttttct	gtgactgggt	agtactcaac	caagtcattc	tgagaatacc	3240
gcgcccggcg	accgagttgc	tcttgcccgg	cgtcaatacg	ggataatagt	gtatgacata	3300
gcagaacttt	aaaagtgtc	atcattggaa	aacgttcttc	ggggcgaaaa	ctctcaagga	3360
tcttaccgct	gttgagatcc	agttcgatgt	aaccactcgc	tgacccaac	tgatcttcag	3420
catcttttac	tttcaccagc	gtttctgggt	gagcaaaaac	aggaaggcaa	aatgccgcaa	3480
aaaagggaat	aaggcgacac	cggaaatgtt	gaatactcat	actcttcctt	tttcaatatt	3540
attgaagcat	ttatcagggg	tattgtctca	tgagcggata	catatttgaa	tgtatttaga	3600
aaaataaaca	aataggggtt	ccgcgcacat	ttccccgaaa	agtgccacct	gacgtctaag	3660
aaaccattat	tatcatgaca	ttaacctata	aaaataggcg	tatcacgagg	ccctttcgtc	3720
tcgcgcggtt	cggtagtgac	ggtgaaaacc	tctgacacat	gcagctcccg	gagacgggtc	3780
cagcttgtct	gtaagcggat	gccgggagca	gacaagcccg	tcagggcgcg	tcagcgggtg	3840
ttggcgggtg	tcggggctgg	cttaactatg	cggcatcaga	gcagattgta	ctgagagtgc	3900
accatatgcg	gtgtgaaata	ccgcacagat	gcgtaaggag	aaaataccgc	atcaggcgaa	3960
attgtaaacg	ttaatathtt	gttaaaattc	gcgttaata	tttgttaaat	cagctcattt	4020
tttaaccaat	aggccgaaat	cggcaaaatc	ccttataaat	caaaagaata	gaccgagata	4080
gggttgagtg	ttgttccagt	ttggaacaag	agtccactat	taaagaacgt	ggactccaac	4140
gtcaaagggc	gaaaaaccgt	ctatcagggc	gatggcccac	tacgtgaacc	atcacccaaa	4200
tcaagttttt	tgcggtcgag	gtgccgtaaa	gctctaaatc	ggaaccctaa	agggagcccc	4260
cgatttagag	cttgacgggg	aaagccggcg	aacgtggcga	gaaaggaagg	gaagaaagcg	4320
aaaggagcgg	gcgctagggc	gctggcaagt	gtagcgggtc	cgctgcgcgt	aaccaccaca	4380
cccgcgcgc	ttaatgcgcc	gctacagggc	gcgtccattc	gccattcagg	ctgcgcaact	4440
gttggaagg	gcgatcggtg	cgggcctctt	cgctattacg	ccagctggcg	aaagggggat	4500
gtgctgcaag	gcgattaagt	tgggtaacgc	cagggttttc	ccagtcacga	cgttgtaaaa	4560
cgacggccag	tgaattgtaa	tacgactcac	tatagggcga	attcgagctc	ggtaccgggg	4620

&lt;210&gt; 3

&lt;211&gt; 4756

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Plasmid pGX22

&lt;400&gt; 3

tgctcagaga	gtttctcaac	gaacccgatt	tggctagtta	taggtaattt	ttagaacatt	60
tacaaaaaca	gcaaaaaaac	caaacattca	ggatttttgt	ttttaattaa	gaaaaaaatc	120
gatcgctctt	aaattttaat	caatacttcg	aataaaccca	aaaaaaaacg	aaaaaaaatc	180
ctgtttccag	tgtaatgatg	attgacgagg	ctcacgaacg	tactctacac	acggatattc	240
tattcaggtt	agtcaaagat	attgcaagat	tccgaaagga	tttgaagctt	ctcatctctt	300
ctgcaacact	tgacgctgaa	aagttctcca	gtttcttcga	cgacgctccg	attttccgaa	360
ttccgggaog	cagattcccc	gtggacattt	actatacaca	ggctcccgaa	gcggactacg	420
tcgacgcggc	tatcgtcaca	attatgcaga	ttcacttgac	ccagccactt	cccggcgata	480
ttttgggtatt	tctgactggt	caggaagaaa	tcgaaactgt	acaggaagca	cttatggaac	540
ggtcgaaagc	actgggatcg	aagattaagg	agcttattcc	gctgccggtt	tatgcgaatt	600
tgcccagtg	tttgaggcg	aagattttcg	agccaacgcc	gaaagatgcg	agaaaggtag	660
atttttctta	caaatttttt	ccaaaaaaa	atccgagaaa	aatctacaaa	atttcaggca	720
aaaactgttt	catttttattc	ctaactagtt	tttttagcaa	cgtttagatt	taacaaaact	780
gaacaaattt	gaagttttcc	aattttaaaaa	ataaatgttt	cggaaaagttt	attgaaaaat	840
ctgaaattgc	tatcctctcg	tatctgcaaa	aaaaacactt	taaaaaatgc	tctgttcttt	900
gaaaattttct	aaactgaaaa	atttgaaatt	tctgaaaatt	gtgataattt	tataaaaatt	960
tatagaaaat	gtaagcattc	cagaaaaata	tcaaaaaatt	cgagaaaaatt	ctgaaaaaat	1020
ccagaaatat	taacagaaaa	aaaatctttt	gaaacatctg	aaaatttaaaa	taaattgaat	1080
ttacattttt	ttttttggga	tttctttaa	atcactatga	atttaccact	aaattttttg	1140
caaaaaatta	tttttttaaat	ttcaaagaaa	aagcaaagaa	ttttaaaata	tcaaaaagtc	1200
caaatttggt	tcggtgaatt	tttaaaataa	cattttcaag	ataattttta	gttaatcaaa	1260
acattccacg	cattttctagt	ttcccaaat	tctctaaatt	tcaggtggtc	ctagcaacta	1320
acattgccag	cacaatggat	ctcgagggat	cttccatacc	taccagttct	gcgcctgcag	1380
gtcgcggccg	cgactctcta	gacgcgtaag	cttactagca	taacccttg	gggctcttaa	1440
acgggtcttg	aggggttttt	tgagcttctc	gccctatagt	gagtcgtatt	acagcttgag	1500
tattctatag	tgtcacctaa	atagcttgcc	gtaactatgg	tcatagctgt	tcctgtgtgt	1560
aaattgttat	ccgctcacaa	ttccacacaa	catagcagcc	ggaagcataa	agtgtaaaagc	1620
ctggggtgcc	taatgagtga	gctaactcac	attaattgcy	ttgcgctcac	tgcccgcctt	1680
ccagtcggga	aacctgtcgt	gccagctgca	ttaatgaatc	ggccaacgcg	cggggagagg	1740
cggtttgcgt	attgggcgct	cttcgccttc	ctcgcctact	gactcgctgc	gctcggctcg	1800
tcggctgcgg	cgagcgggat	cagctcactc	aaaggcggta	atacggttat	ccacagaatc	1860
aggggataac	gcaggaaaga	acatgtgagc	aaaaggccag	caaaaggcca	ggaaccgtaa	1920
aaaggccgcy	ttgctggcgt	ttttcgatag	gctccgcccc	cctgacgagc	atcacaaaaa	1980
tcgacgctca	agtcagaggt	ggcgaaaccc	gacaggacta	taaagatacc	aggcgttttc	2040
ccctggaagc	tccctcgtgc	gctctcctgt	tccgaccctg	ccgcttaccg	gatacctgtc	2100
cgctttcttc	ccttcgggaa	gcgtggcgct	ttctcatagc	tcacgctgta	ggtatctcag	2160
ttcgggtgtag	gtcgttcgct	ccaagctggg	ctgtgtgcac	gaaccccccg	ttcagcccga	2220
ccgctgcgcc	ttatccggta	actatcgtct	tgagtccaac	ccggtaagac	acgacttatc	2280
gccactggca	gcagccactg	gtaacaggat	tagcagagcg	aggatgttag	gcggtgctac	2340
agagttcttg	aagtgggtggc	ctaactacgg	ctacactaga	aggacagtat	ttggtatctg	2400
cgctctgctg	aagccagtta	ccttcggaaa	aagagttggt	agctcttgat	ccggcaaaaa	2460
aaccaccgct	ggtagcgggt	gtttttttgt	ttgcaagcag	cagattacgc	gcagaaaaaa	2520
aggatctcaa	gaagatcctt	tgatcttttc	tacggggtct	gacgctcagt	ggaacgaaaa	2580
ctcacgttaa	gggatttttg	tcatgagatt	atcaaaaagg	atcttcacct	agatcctttt	2640
aaattaaaaa	tgaagtttta	aatcaatcta	aagtatatat	gagtaaaact	ggtctgacag	2700
ttaccaatgc	ttaatcagtg	aggcacctat	ctcagcgatc	tgtctatttc	gttcatccat	2760
agttgcctga	ctccccgtcg	tgtagataac	tacgatacgg	gagggcttac	catctggccc	2820
cagtgtcgca	atgataccgc	gagacccacg	ctcaccggct	ccagatttat	cagcaataaa	2880
ccagccagcc	ggaagggccg	agcgcagaag	tggtcctgca	actttatccg	cctccatcca	2940
gtctattaat	tggtgccggg	aagctagagt	aagtagttcg	ccagttaata	gtttgcgcaa	3000
cgttgttggc	attgctacag	gcacgtggt	gtcacgctcg	tcgtttggt	tggcttcatt	3060
cagctccggt	tcccaacgat	caaggcgagt	tacatgatcc	cccatgttgt	gcaaaaaagc	3120
ggttagctcc	ttcggtcctc	cgatcgttgt	cagaagtaag	ttggccgcag	tgttatcact	3180
catggttatg	gcagcactgc	ataattctct	tactgtcatg	ccatccgtaa	gatgcttttc	3240
tgtgactggt	gagtactcaa	ccaagtcatt	ctgagaatac	cgcgcccgcc	gaccgagttg	3300
ctcttgcccc	gcgtcaatac	gggataatag	tgtatgacat	agcagaactt	taaaagtgc	3360

```

catcattgga aaacgttctt cggggcgaaa actctcaagg atcttaccgc tgttgagatc 3420
cagttcgaatg taaccactc gtgcacccaa ctgatcttca gcatctttta ctttcaccag 3480
cgtttctggg tgagcaaaaa caggaaggca aaatgccgca aaaaagggaa taaggcgac 3540
acggaaatgt tgaatactca tactcttcct ttttcaatat tattgaagca tttatcaggg 3600
ttattgtctc atgagcggat acatatttga atgtatttag aaaaataaac aaataggggt 3660
tccgcgcaca tttccccgaa aagtgccacc tgacgtctaa gaaaccatta ttatcatgac 3720
attaacctat aaaaataggc gtatcacgag gccctttcgt ctgcgcggtt tcggtgatga 3780
cggtgaaaaac ctctgacaca tgcagctccc ggagacgggt acagcttgtc tgtaagcggg 3840
tgccgggagc agacaagccc gtcagggcgc gtcagcgggt gttggcgggt gtcggggctg 3900
gcttaactat gcggcatcag agcagattgt actgagagtg caccatatgc ggtgtgaaat 3960
accgcacaga tgcgtaaggg gaaaataacc catcaggcga aattgtaaac gttaatat 4020
tgtaaaatt cgcggttaaat atttgttaaa tcagctcatt ttttaaccaa taggccgaaa 4080
tcggcaaaat cccttataaa tcaaaagaat agaccgagat aggggtgagt gttgttccag 4140
tttggaacaa gagtccacta ttaaagaacg tggactccaa cgtcaaaggg cgaaaaaccg 4200
tctatcaggg cgatggccca ctacgtgaac catcacccaa atcaagtttt ttgcggtcga 4260
ggtgccgtaa agctctaaat cggaacccta aaggggagccc ccgatttaga gcttgacggg 4320
gaaagccggc gaacgtggcg agaaaggaag ggaagaaagc gaaaggagcg ggcgctaggg 4380
cgctggcaag tgtagcgggt acgctgcgcg taaccaccac acccgccgcg cttaatgcgc 4440
cgctacaggg cgcgctccatt cgccattcag gctgcgcaac tgttgggaag ggcgatcgg 4500
gcgggcctct tcgctattac gccagctggc gaaaggggga tgtgctgcaa ggcgattaa 4560
ttgggtaacg ccagggtttt cccagtcacg acgttgtaaa acgacggcca gtgaattgta 4620
atacgactca ctatagggcg aattcaaaaa acccctcaag acccgtttag aggcccaag 4680
gggttatgct agtgaattct gcagggtacc cggggatcct ctagagatcc ctcgacctcg 4740
agatccattg tgctgg 4756

```

&lt;210&gt; 4

&lt;211&gt; 4643

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Plasmid pGX52

&lt;400&gt; 4

```

gagtgcacca tatgcggtgt gaaataccgc acagatgcgt aaggagaaaa taccgcatca 60
ggcgaaattg taaacgttaa tattttgtaa aaattcgcgt taaatatattg ttaaaccagc 120
tcatttttta accaataggc cgaaatcggc aaaatccctt ataaatcaaa agaataagacc 180
gagatagggt tgagtgttgt tccagtttgg aacaagagtc cactattaaa gaacgtggac 240
tccaacgtca aagggcgaaa aaccgtctat cagggcgatg gccactacg tgaaccatca 300
cccaaatcaa gttttttgcg gtcgaggtgc cgtaaagctc taaatcgga ccctaaagg 360
agccccgat ttagagcttg acggggaaa cggcggaacg tggcgagaaa ggaagggaa 420
aaagcgaaag gagcgggcgc tagggcgctg gcaagtgtag cggtcacgct gcgcgtaacc 480
accacaccgg ccgcgcttaa tgcgcgcta cagggcgctt ccattcgcca ttcaggctgc 540
gcaactgttg ggaagggcga tcggtgcggg cctcttcgct attacgccag ctggcgaaa 600
ggggatgtgc tgcaaggcga ttaagtggg taacgccagg gttttcccag tcacgacgtt 660
gtaaaacgac ggccagtga ttgtaatacg actcactata gggcgaaattc aaaaaacccc 720
tcaagaccgg ttagaggcc ccaaggggtt atgctagtga attctgcagg gtaccgggg 780
atcctctaga gatccctcga cctcgagatc cattgtgctg gcagccgatc tccgtctgt 840
gaagatctac tccaccacca tccgtatcga tcagtccatc ctcaccggag aatctgtgtc 900
tgttatcaag cacaccgact ctgtgccaga tccacgcgct gttaaccagg acaagaagaa 960
ttgtctgttc tcgggaacca atgtcgcac tggaagggt cgtggaatcg tcttcggaac 1020
cggattgacc actgaaatcg gaaagatccg taccgaaatg gctgagaccg agaatgagaa 1080
gacaccactt caacagaagt tggacgaatt cggagagcaa ctttccaagg ttatctctgt 1140
tatgtcggtt gctgtttggg ctatcaacat tggacatttc aacgatccag ctcacgggtg 1200
atcatgggtt aaggggagcaa tctactactt caaaatcgcc gttgctcttg ccgtcgctgc 1260
tattccagaa ggacttccag ctgtcatcac cacgtgcctt gccctcgga ctcgccgtat 1320
ggccaagaag aacgctattg taagatccct tccatccgtc gaaactcttg gatgcacatc 1380
tgttatctgc tctgacaaga ctggaactct caccaccaac cagatgtctg tgtcaaagat 1440

```

gttcatcgct	ggacaagctt	ctggagacaa	catcaacttc	accgagttcg	ccatctccgg	1500
atccacctac	gagccagtcg	gaaaggtttc	caccaatgga	cgtgaaatca	acccagctgc	1560
tggagaattc	gaatcactca	ccgagttggc	catgatctgc	gctatgtgca	atgattcatc	1620
tggttgattac	aatgagacca	agaagatcta	cgagaaagtc	ggagaagcca	ctgaaactgc	1680
tcttatcggt	cttgctgaga	agatgaatgt	tttcggaacc	tcgaaagccg	gactttcacc	1740
aaaggagctc	ggaggagttt	gcaaccgtgt	catccaacaa	aaatggaaga	aggagttcac	1800
actcgagttc	tcccgtgata	gtaaattccat	gtccgcctac	tgcttcccag	cttccggagg	1860
atctggagcc	aagatgttcg	tgaaggggagc	cccagaagga	gttctcggaa	gatgcacca	1920
cgtcagagtt	aacggacaaa	aggttccact	cacctctgcc	atgactcaga	agattgttga	1980
ccaatgcgtg	caatacggaa	ccggaagaga	tacccttcgt	tgtcttgccc	tcggccagca	2040
caatggatct	cgagggatct	tccataccta	ccagttctgc	gcctgcaggt	cgcgcccgcg	2100
actctctaga	cgcgtaagct	tactagcata	accccttggg	gcctctaaac	gggtcttgag	2160
gggttttttg	agcttctcgc	cctatagtga	gtcgtattac	agcttgagta	ttctatagt	2220
tcacctaaat	agcttggcgt	aatcatggtc	atagctgttt	cctgtgtgaa	attgttatcc	2280
gctcacaatt	ccacacaaca	tacgagccgg	aagcataaag	tgtaaagcct	ggggtgccta	2340
atgagtgage	taactcacat	taattgcgtt	gcgctcactg	cccgttttcc	agtcgggaaa	2400
cctgtcgtgc	cagctgcatt	aatgaatcgg	ccaacgcgcg	gggagaggcg	gtttgcgtat	2460
tgggcgctct	tccgcttcc	cgctcactga	ctcgtgcgc	tcggtcgttc	ggctgcggcg	2520
agcggtatca	gtcactcaa	aggcggtaat	acggttatcc	acagaatcag	gggataacgc	2580
aggaaagaac	atgtgagcaa	aaggccagca	aaaggccagg	aaccgtaaaa	aggccgcgtt	2640
gctggcggtt	ttcgataggc	tccgcccccc	tgacgagcat	cacaaaaatc	cagcgtcaag	2700
tcagaggtgg	cgaaacccga	caggactata	aagataccag	gcgtttcccc	ctggaagctc	2760
cctcgtgcgc	tctcctgttc	cgacctgcc	gcttaccgga	tacctgtccg	cctttctccc	2820
ttcgggaagc	gtggcgcttt	ctcatagctc	acgctgtagg	tatctcagtt	cggtgtaggt	2880
cgttcgctoc	aagctgggct	gtgtgcacga	acccccggt	cagcccagacc	gctgcgcctt	2940
atccggtaac	tatcgtcttg	agtccaaccc	ggtaagacac	gacttatcgc	cactggcagc	3000
agccactgg	aacaggatta	gcagagcgag	gtatgtaggc	ggtgctacag	agttcttgaa	3060
gtggtggcct	aactacggct	acactagaag	gacagtattt	ggtatctgcg	ctctgctgaa	3120
gccagttacc	ttcggaaaaa	gagttggtag	ctcttgatcc	ggcaaacaaa	ccaccgctgg	3180
tagcggtggt	ttttttgttt	gcaagcagca	gattacgcgc	agaaaaaaag	gatctcaaga	3240
agatccattg	atcttttcta	cggggtctga	cgctcagtg	aacgaaaact	cacgttaagg	3300
gattttggtc	atgagattat	caaaaaggat	cttcacctag	atccttttaa	attaaaaatg	3360
aagtttttaa	tcaatctaaa	gtatatatga	gtaaaacttg	tctgacagtt	accaatgctt	3420
aatcagtgag	gcacctatct	cagcgatctg	tctatttctg	tcattccatg	ttgcctgact	3480
ccccgctcgt	tagataacta	cgatacggga	gggcttacca	tctggcccca	gtgctgcaat	3540
gataccgcga	gacccacgct	caccggctcc	agatttatca	gcaataaacc	agccagccgg	3600
aaggggccgag	cgcagaagtg	gtcctgcaac	tttatccgcc	tccatccagt	ctattaattg	3660
ttgccgggaa	gctagagtaa	gtagtctgcc	agttaatagt	ttgcgcaacg	ttgttggtac	3720
tgctacaggc	atcgtggtgt	cacgctcgtc	gtttggtatg	gcttcattca	gctccggttc	3780
ccaacgatca	aggcgagtta	catgatcccc	catgttgtgc	aaaaaaagcg	ttagctcctt	3840
cggctcctccg	atcgttgtca	gaagtaagtt	ggccgcagtg	ttatcactca	tggttatggc	3900
agcactgcat	aattctctta	ctgtcatgcc	atccgtaaga	tgcttttctg	tgactggtga	3960
gtactcaacc	aagtcattct	gagaataaccg	cgccccggcg	ccgagttgct	cttgcccggc	4020
gtcaatacgg	gataatagtg	tatgacatag	cagaacttta	aaagtgtctc	tcattggaaa	4080
acgttcttcg	gggcgaaaaa	tctcaaggat	cttaccgctg	ttgagatcca	gttcgatgta	4140
acccactcgt	gcacccaact	gatcttcagc	atcttttact	ttcaccagcg	tttctgggtg	4200
agcaaaaaa	ggaaggcaaa	atgccgcaaa	aaagggaata	agggcgacac	ggaaatgttg	4260
aatactcata	ctcttccctt	ttcaatatta	ttgaagcatt	tatcagggtt	attgtctcat	4320
gagcggatac	atatttgaat	gtatttagaa	aaataaacia	ataggggttc	cgcgcacatt	4380
tccccgaaaa	gtgccacctg	acgtctaaga	aaccattatt	atcatgacat	taacctataa	4440
aaataggcgt	atcacgaggc	cctttcgtct	cgcgcggttc	ggtgatgacg	gtgaaaacct	4500
ctgacacatg	cagctcccgg	agacggtcac	agcttgtctg	taagcggatg	ccgggagcag	4560
acaagcccgt	cagggcgcgt	cagcgggtgt	tggcgggtgt	cggggctggc	ttactatgc	4620
ggcatcagag	cagattgtac	tga				4643

&lt;210&gt; 5

&lt;211&gt; 4454

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Plasmid pGX104

&lt;400&gt; 5

```

gagtgcacca tatgcggtgt gaaataccgc acagatgcgt aaggagaaaa taccgcatca 60
ggcgaaaattg taaacgttaa tattttgtta aaattcgcgt taaatatttg ttaaatacagc 120
tcattttttta accaataggc cgaaatcggc aaaatccctt ataaatcaaa agaatagacc 180
gagatagggt tgagtgttgt tccagtttg aacaagagtc cactattaaa gaacgtggac 240
tccaacgtca aaggcgaaaa aaccgtctat cagggcgatg gccactacg tgaaccatca 300
cccaaatcaa gttttttgcg gtcgaggtgc cgtaaagctc taaatcggaa ccctaaagg 360
agccccgat ttagagcttg acggggaaa cggcgaaacg tggcgagaaa ggaagggaag 420
aaagcgaaag gagcgggccc tagggcgctg gcaagtgtag cggtcacgct gcgcgtaacc 480
accacacccg ccgcgcttaa tgcgcccgtc cagggcgctt ccattcgcca ttcaggctgc 540
gcaactgttg ggaaggcgga tccgtgcggg cctcttcgct attacgccag ctggcgaaa 600
ggggatgtgc tgcaaggcga ttaagtggg taacgccagg gttttcccag tcacgacgtt 660
gtaaacgcag ggccagtga ttgtaatac actcactata gggcgaattc aaaaaacccc 720
tcaagaccg ttttagagg ccaagggggt atgctagtga attctgcagg gtacccggg 780
atcctctaga gatccctcga cctcgagatc cattgtgctg gaccgtggta ctcttatgga 840
gctcggaatc tgcgaatcg tcaattctg acttatcatg caacttctc cggagccaa 900
gatcatcgaa gtcggagaca caccaaagga ccgtgctctt ttcaacggag ccagaaatg 960
taagccgaaa agtgtgtgtt ttcaatctct aatttttgaa cttttcagtg ttcggtatgg 1020
tcatcactgt tggacaagct attgtctacg tcatgtccgg actctacgga gagccatcg 1080
aaatcggagc tggaaatctgt ctccattatc tctccaact cgttattgcc ggtctcatcg 1140
tcctccttct cgacgagctt ctccaaaagg gatattggtc cggatccgga atttctctct 1200
tcattgccac caacatctgt gaaaccattg tctggaaggc attctccccg gcaacaatga 1260
acacggagc tggaaccgag ttogaaggag ccgtcattgc tcttttccat ctcttgcca 1320
cccgctccga caaggtccgt gcccttcgtg aggccttcta ccgtcaaaac cttccaaact 1380
tgatgaactt gatggctact ttctcgttt ttgcggtggg tatctacttc caaggattcc 1440
gtgtcgacct cccaatcaag tctgcccgt accgtggaca atacagcagc taccaatca 1500
agctcttcta cacctccaac attccaatca tcttcaatc tgctctcgtc tccaacctct 1560
acgttatctc tcaggtttgt tgcattctcag tagtaccgtt agatgtttat ctttctctag 1620
agggtcaggt tggcgagaaa attttttgag ttcattctca agtctgatgg aaaatgttta 1680
tttttcagat gtcgcccggg aagttcggag gaaacttctt catcaacctt ctcggtacct 1740
ggtccgataa caccggatag agaagctacc caactggagg actctgctac tatctttcac 1800
caccagagtc tcttgacac atcttcgaag acccaatcca ctgcaccagc acaatggagt 1860
tcgagggagc ttccatacct accagttctg cgcctgcagg tgcgcccgcg gactctctag 1920
acgcgtaagc ttactagcat aacccttggt ggccctctaaa cgggtcttga ggggtttttt 1980
gagcttctcg ccctatagtg agtcgtatta cagcttgagt attctatagt gtcacctaaa 2040
tagcttggcg taatcatggt catagctgtt tctgtgtga aattgttatc cgctcacaat 2100
tccacacaac atacgagccg gaagcataaa gtgtaaagcc tggggtgcct aatgagttag 2160
ctaactcaca ttaattgcgt tgcgctcact gccgcgttcc cagtcgggaa acctgtcgtg 2220
ccagctgcat taatgaatcg gccaacgcgc ggggagaggc ggtttgcgta ttgggcgctc 2280
ttccgcttcc tcgctcactg actcgtgcg ctcggtcgtt cggctgcggc gagcggtatc 2340
agctcactca aaggcggtaa tacggttatc cacagaatca ggggataacg caggaaagaa 2400
catgtgagca aaaggccagc aaaaggccag gaaccgtaaa aaggccgcgt tgctggcggt 2460
tttcgatagg ctccgcccc ctgacgagca tcacaaaaat cgacgctcaa gtcagagggt 2520
gcgaaacccg acaggactat aaagatacca ggcgtttccc cctggaagct ccctcgtgcg 2580
ctctcctgtt ccgacctgc cgcttaccgg atacctgtcc gcctttctcc cttcggaag 2640
cgtggcgctt tctcatagct cacgctgtag gtatctcagt tcggtgtagg tcgttcgctc 2700
caagctgggc tgtgtgcacg aacccccgt tcagcccagc cgctgcgcct tatccggtaa 2760
ctatcgtctt gagtccaacc cgtaagaca cgacttatcg ccactggcag cagccactgg 2820
taacaggatt agcagagcga ggtatgtagg cgggtctaca gagttcttga agtgggtggc 2880
taactacggc tacactagaa ggacagtatt tggatatctg gctctgctga agccagttac 2940
cttcggaaaa agagttggta gctcttgatc cggcaaaaa accaccgctg gtacgggtgg 3000
tttttttgtt tgcaagcagc agattacgcg cagaaaaaaa ggatctcaag aagatcctt 3060
gatcttttct acgggggtctg acgctcagtg gaacgaaaa tcacgttaag ggattttggt 3120

```



```

catgagatta tcaaaaagga tcttcaccta gatcctttta aattaaaaat gaagttttta 3180
atcaatctaa agtatatatg agtaaaactg gtctgacagt taccaatgct taatcagtga 3240
ggcacctatc tcagcgatct gtctatctcg ttcacccata gttgcctgac tccccgctcg 3300
gtagataact acgatacggg agggccttacc atctggcccc agtgctgcaa tgataccgcg 3360
agaccacgc tcaccggctc cagatttatc agcaataaac cagccagccg gaagggccga 3420
gcgcagaagt ggctctgcaa ctttatccgc ctccatccag tctattaatt gttgccggga 3480
agctagagta agtagttcgc cagttaatag tttgcgcaac gttgttgga ttgctacagg 3540
catcgtggtg tcacgctcgt cgtttggtat ggcttcattc agctccggtt cccaacgac 3600
aaggcgagtt acatgatccc ccatgtttgtg caaaaaagcg gttagctcct tcggtcctcc 3660
gatcgtttgc agaagtaagt tggccgcagt gttatcactc atgggttatg cagcactgca 3720
taattctctt actgtcatgc catccgtaag atgcttttct gtgactggtg agtactcaac 3780
caagtcatte tgagaatacc gcgcccggcg accgagttgc tcttgcccgg cgtcaatacg 3840
ggataatagt gtatgacata gcagaacttt aaaagtgtc atcattggaa aacgttcttc 3900
ggggcgaaaa ctctcaagga tcttaccgct gttgagatcc agttcgatgt aaccactcg 3960
tgcacccaac tgatcttcag catcttttac tttcaccagc gtttctgggt gagcaaaaac 4020
aggaaggcaa aatgccgcaa aaaaggggaat aagggcgaca cggaaatgtt gaatactcat 4080
actcttctt tttcaatatt attgaagcat ttatcagggt tattgtctca tgagcggata 4140
catatttgaa tgtatttaga aaaataaaca aataggggtt ccgcgcacat ttccccgaaa 4200
agtgccacct gacgtctaag aaaccattat tatcatgaca ttaacctata aaaataggcg 4260
tatcacgagg ccttttcgtc tcgcgcgttt cggtagatgc ggtgaaaacc tctgacacat 4320
gcagctcccg gagacggtca cagcttgtct gtaagcggtat gccgggagca gacaagcccg 4380
tcagggcgcg tcagcgggtg ttggcgggtg tcggggctgg cttaactatg cggcatcaga 4440
gcagattgta ctga 4454

```

&lt;210&gt; 6

&lt;211&gt; 4701

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Plasmid pGZ18

&lt;400&gt; 6

```

accagcttt cttgtacaaa gtggtgatct ttccagcaca atggatctcg agggatcttc 60
catacctacc agttctgcgc ctgcaggtcg cggccgcgac tctctagacg cgtaagctta 120
ctagcataac ccttgggggc ctctaaacgg gtcttgaggg gttttttgag cttctcgccc 180
tatagtgagt cgtattacag cttgagtatt ctatagtgtc acctaaatag cttggcgtaa 240
tcatggtcat agctgtttcc tgtgtgaaat tggtatccgc tcacaattcc acacaacata 300
cgagccggaa gcataaagtg taaagcctgg ggtgcctaag gtagtgagta actcacatta 360
attgcgttgc gctcactgcc cgctttccag tcgggaaacc tgtcgtgccca gctgcattaa 420
tgaatcggcc aacgcgcggg gagaggcggg ttgcgtattg ggcgctcttc cgcttcctcg 480
ctcactgact cgctgcgctc ggtcgttcgg ctgcggcgag cggtatcagc tcaactaaag 540
gcggtataac ggttatccac agaatacagg gataacgcag gaaagaacat gtgagcaaaa 600
ggccagcaaa aggccaggaa ccgtaaaaag gccgcgttgc tggcgttttt cgataggctc 660
cgccccctg acgagcatca caaaaatcga cgctcaagtc agaggtggcg aaaccgcaga 720
ggactataaa gataaccaggc gtttccccct ggaagctccc tcgtgcgctc tcctgttccg 780
accctgccgc ttaccggata cctgtccgcc tttctccctt cgggaagcgt ggcgctttct 840
catagctcac gctgtaggta tctcagttcg gtgtaggctg ttcgctccaa gctgggctgt 900
gtgcacgaac cccccgttca gcccgaccgc tgcgccttat ccggtactta tcgtcttgag 960
tccaaccogg taagacacga cttatcgcca ctggcgagcag ccactggtaa caggattagc 1020
agagcgaggt atgtaggcgg tgctacagag ttcttgaagt ggtggcctaa ctacggctac 1080
actagaagga cagtatttgg tatctgcgct ctgctgaagc cagttacctt cggaaaaaga 1140
gttggttagct cttgatccgg caaacaacc accgctggta gcggtggttt ttttgtttgc 1200
aagcagcaga ttacgcgcag aaaaaagga tctcaagaag atcctttgat cttttctacg 1260
gggtctgacg ctacgtggaa cgaaaactca cgtaaggga ttttgggtcat gagattatca 1320
aaaaggatct tcacctagat ctttttaaat taaaaatgaa gttttaaatc aatctaaagt 1380
atatatgagt aaacttggtc tgacagttac caatgcttaa tcagtgaggc acctatctca 1440
gcgatctgtc tatttcgttc atccatagtt gcctgactcc ccgtcgtgta gataactacg 1500

```

atacgggagg	gottaccatc	tgccccag	gctgcaatga	taccgcgaga	cccacgctca	1560
ccggctccag	atztatcagc	aataaaccag	ccagccggaa	gggcccagcg	cagaagtgg	1620
cctgcaactt	tatccgcctc	catccagctc	attaattggt	gccgggaagc	tagagtaagt	1680
agttcgccag	ttaatagttt	gcgcaacggt	gttggcattg	ctacaggcat	cgtgggtgtca	1740
cgctcgctgt	ttggtatggc	ttcattcagc	tccgggtccc	aacgatcaag	gcgagttaca	1800
tgatccccc	tgttgtgcaa	aaaagcggtt	agctccttcg	gtcctccgat	cgttgtcaga	1860
agtaagttgg	ccgcagtggt	atcactcatg	gttatggcag	cactgcataa	ttctcttact	1920
gtcatgccat	ccgtaagatg	cttttctgtg	actggtgagt	actcaaccaa	gtcattctga	1980
gaataccgag	ccggcgacc	gagttgctct	tgcccggcgt	caatacggga	taatagtgtg	2040
tgacatagca	gaactttaaa	agtgtctatc	attggaaaac	gttcttcggg	gcgaaaactc	2100
tcaaggatct	taccgctggt	gagatccagt	tcgatgtaac	ccactcgtgc	acccaactga	2160
tcttcagcat	cttttacttt	caccagcggt	tctgggtgag	caaaaacagg	aaggcaaaat	2220
gccgcaaaaa	aggggaataag	ggcgacacgg	aaatgttgaa	tactcatact	cttccttttt	2280
caatattatt	gaagcattta	tcagggttat	tgtctcatga	gcggatacat	atttgaatgt	2340
atttagaaaa	ataaacaat	aggggttccg	cgcacatttc	cccgaagaag	gccacctgac	2400
gtctaagaaa	ccattattat	catgacatta	acctataaaa	ataggcggtat	cacgaggccc	2460
tttcgtctcg	cgcggttcgg	tgatgacggt	gaaaacctct	gacacatgca	gctcccggag	2520
acggtcacag	cttgtctgta	agcggatgcc	gggagcagac	aagcccgtca	gggcgctca	2580
gcgggtgttg	ccgggtgtcg	gggctggctt	aactatgcgg	catcagagca	gattgtactg	2640
agagtgcacc	atatgcgggt	tgaaataccg	cacagatgcg	taaggagaaa	ataccgcattc	2700
aggcgaaatt	gtaaacgtta	atattttgtt	aaaattcgcg	ttaaatattt	gttaaatcag	2760
ctcatttttt	aaccaatagg	ccgaaatcgg	caaaatccct	tataaatcaa	aagaatagac	2820
cgagataggg	ttgagtgttg	ttccagtttg	gaacaagagt	ccactattaa	agaacgtgga	2880
ctccaacgtc	aaagggcgaa	aaaccgtcta	tcaggggcat	ggcccactac	gtgaaccatc	2940
acccaaatca	agttttttgc	ggtcgagggtg	ccgtaaagct	ctaaatcgga	accctaaagg	3000
gagcccccca	tttagagctt	gacggggaaa	gccggcgaa	gtggcgagaa	aggaagggaa	3060
gaaagcgaaa	ggagcgggcg	ctagggcgct	ggcaagtgtg	gcggtcacgc	tgccgctaac	3120
caccacaccc	gccgcgctta	atgcgcgcgt	acagggcgcg	tccattcgcc	attcaggctg	3180
cgcaactggt	gggaaggcg	atcggtgcgg	gcctcttcgc	tattacgcca	gctggcgaaa	3240
gggggatgtg	ttgcaaggcg	attaagttgg	gtaacgccag	ggttttccca	gtcacgacgt	3300
tgtaaaacga	cggccagtg	attgtaatac	gactcactat	agggcggaat	caaaaaaccc	3360
ctcaagaccc	gttttagagg	cccaaggggt	tatgctagt	aattctgcag	ggtaccggg	3420
gatcctctag	agatccctcg	acctcgagat	ccattgtgct	ggaaagcctt	tgagggtcg	3480
gcaagccacg	tttggtggtg	gcgaccatcc	tccaaaatca	acaagtgtgt	acaaaaaagc	3540
aggctatgcc	aagtacatgt	cgattgcgta	cgcgttcgta	atgttggctg	tgtagtcgc	3600
taccagcagt	caaattgttc	tcgagagtgc	gtttttacat	tatcccttca	tcctgattac	3660
gacaattttc	agctgttctc	gtcctctcat	ctctcttcat	tgtcacaatg	gtcggaatct	3720
tcttctttgc	tgattgtctt	catccaaaag	aattcacgaa	tattatccat	ggtgtcgat	3780
tcttctctcat	gattccatct	acatatgtgt	tctcactttt	atattcgctc	atcaatctca	3840
acgttatcac	gtggggaaact	cgtgaagctg	tcgctaaggc	aacgggacaa	aagacgaaaa	3900
aagcgccctat	ggaacaattt	atagacagag	tgattgatat	tgtgaaaaag	ggattcagat	3960
taatcagttg	tcggggagaag	aaggaacatg	aagagagacg	agagaaaatg	gaaaagaaaa	4020
tgagagaaat	ggagctagcc	ttgagaagta	ttgagggttat	ctttaacttt	agaaatgtga	4080
aattaataat	ttatttttcag	agtgggtgccg	acgtgaagaa	aattctcgat	gcaacagagg	4140
agaaggagaa	acgtgaagaa	gaaactcaaa	ctgcagattt	tccgattgaa	gagaacgtag	4200
agaagactca	aaaagagatt	cagaaggcaa	accgttatgt	gtggatgaca	agtcatagct	4260
tgaagtttg	tgaacgagga	aaactgaaaa	gtgcggaaaa	ggttttctgg	aacgagctca	4320
tcaatgcata	tctgaaaccg	atcaagacga	cgcagctga	aatgaaagcc	gtcgccgaag	4380
gattggcttc	tctacgaaat	cagattgctt	tcactattct	tctcgtaaat	tctcttcttg	4440
ctcttgccat	ctttttgatt	cagaaacaca	aaaatgtgct	cagcatcaag	ttctcgccaa	4500
tcagtaagca	atattacctt	tatggtcaat	tcaaaaaatt	tgtttttttt	ttctagaaaa	4560
cttccgatgg	acgaaaatga	atgagatgac	tggaacaatac	gaggaaaccg	atgaaccatt	4620
aaaaatagat	ccacttgga	tggaattgt	tgttttcctt	ctaattatc	ttttgttca	4680
aactctcgga	atgcttctcc	a				4701

&lt;210&gt; 7

&lt;211&gt; 25

&lt;212&gt; DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:  
Oligonucleotide primer C04H5.6F

<400> 7

tgctcagaga gtttctcaac gaacc

25

<210> 8

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Oligonucleotide  
primer C04H5.6R

<400> 8

caatgtagt tgctaggacc acctg

25

<210> 9

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Oligonucleotide  
primer K11D9.2bF

<400> 9

cagccgatct ccgtcttg

20

<210> 10

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Oligonucleotide  
primer K11D9.2bR

<400> 10

ccgagggcaa gacaacgaag

20

<210> 11

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Oligonucleotide  
primer Y57G11C.15F

<400> 11

accgtgtac tcttatggag ctcg

24

<210> 12

<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:Oligonucleotide  
primer Y57G11C.15R

<400> 12  
tgcagtggat tgggtcttcg 20

<210> 13  
<211> 52  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:Oligonucleotide  
primer T25G3.2F

<400> 13  
ggggacaagt ttgtacaaaa aagcaggcta tgccaagtac atgtcgattg cg 52

<210> 14  
<211> 52  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:Oligonucleotide  
primer T25G3.2R

<400> 14  
ggggaccact ttgtacaaga aagctggggtt ggagaagcat tccgagagtt tg 52